



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁵ : C12N 5/00, 15/00, C07H 21/04	A1	(11) International Publication Number: WO 90/06359 (43) International Publication Date: 14 June 1990 (14.06.90)
(21) International Application Number: PCT/US89/05210 (22) International Filing Date: 22 November 1989 (22.11.89) (30) Priority data: 280,218 5 December 1988 (05.12.88) US (71) Applicant: GENPHARM INTERNATIONAL [US/US]; 180 Kimball Way, South San Francisco, CA 94080 (US). (72) Inventors: KRIMPENFORT, Paulus, Jacobus, Angelinus ; Vondelstraat 11 G-IV, NL-1054 GC Amsterdam (NL). BERNS, Antonius, Josef, Maria ; Floris Balthasarstraat 2, NL-2064 XP Spaarndam (NL). (74) Agents: TRECARTIN, Richard, F. et al.; Flehr, Hohbach, Test, Albritton & Herbert, Four Embarcadero Center, Suite 3400, San Francisco, CA 94111-4187 (US).		(81) Designated States: AT (European patent), AU, BE (European patent), CH (European patent), DE (European patent), DK, ES (European patent), FR (European patent), GB (European patent), IT (European patent), JP, LU (European patent), NL (European patent), SE (European patent). Published <i>With international search report.</i>
(54) Title: TRANSGENIC NON-HUMAN ANIMALS DEPLETED IN A MATURE LYMPHOCYTIC CELL-TYPE (57) Abstract Transgenic non-human animals having a phenotype characterized by the substantial depletion of a mature lymphocytic cell type otherwise naturally occurring in the species from which the transgenic animal is derived. The phenotype is conferred in the transgenic animal by a transgene contained in at least the precursor stem cell of the lymphocytic cell type which is depleted. The transgene comprises a DNA sequence encoding a lymphatic polypeptide variant which inhibits maturation of the lymphocytic cell type.		

BEST AVAILABLE COPY

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	ES	Spain	MG	Madagascar
AU	Australia	FI	Finland	ML	Mali
BB	Barbados	FR	France	MR	Mauritania
BE	Belgium	GA	Gabon	MW	Malawi
BF	Burkina Faso	GB	United Kingdom	NL	Netherlands
BG	Bulgaria	HU	Hungary	NO	Norway
BJ	Benin	IT	Italy	RO	Romania
BR	Brazil	JP	Japan	SD	Sudan
CA	Canada	KP	Democratic People's Republic of Korea	SE	Sweden
CF	Central African Republic	KR	Republic of Korea	SN	Senegal
CG	Congo	LI	Liechtenstein	SU	Soviet Union
CH	Switzerland	LK	Sri Lanka	TD	Chad
CM	Cameroon	LU	Luxembourg	TG	Togo
DE	Germany, Federal Republic of	MC	Monaco	US	United States of America
DK	Denmark				

-1-

TRANSGENIC NON-HUMAN ANIMALS DEPLETED
IN A MATURE LYMPHOCYTIC CELL-TYPE

Field of the Invention

The present invention relates to transgenic non-human animals wherein the mature form of at least one lymphocytic cell type is substantially depleted.

5 More particularly, the invention relates to transgenic mice wherein mature T cells or plasma cells are depleted.

Background of the Invention

10 The immune response is a complex defense system that is able to recognize and kill invading organisms such as bacteria, viruses, fungi and possibly also some types of tumor cells. The most characteristic aspects of the immune system are the specific recognition of antigens, the ability to discriminate
15 between self and non-self antigens and a memory-like potential that enables a fast and specific reaction to previously encountered antigens. The vertebrate immune system reacts to foreign antigens with a cascade of molecular and cellular events that
20 ultimately results in the humoral and cell-mediated immune response.

The major pathway of the immune defense commences with the trapping of the antigen by accessory cells such as dendritic cells or macrophages and subsequent concentration in lymphoid organs. There, the accessory cells present the antigen on their cell surface to subclasses of T cells classified as mature T helper cells. Upon specific recognition of the processed antigen the mature T helper cells can be triggered to become activated T helper cells. The activated T helper cells regulate both the humoral immune response by inducing the differentiation of mature B cells to antibody producing plasma cells and control the cell-mediated immune response by activation of mature cytotoxic T cells.

Naturally occurring processes sometimes result in the modulation of immune system cell types. Acquired immunodeficiency syndrome (AIDS) is a devastating infectious disease of the adult immune system which significantly affects cell-mediated immunity. This disease is manifested by profound lymphopenia which appears to be the result of a loss of T-lymphocytes which have the helper/inducer phenotype T4 as defined by the monoclonal antibody OKT4 (Fauci, A., et al. (1984) Annals. Int. Med. 100, 92). Other clinical manifestations include opportunistic infections, predominantly Pneumocystis carinii pneumonia, and Kaposi's sarcoma. Other disease states include Severe Combined Immuno Deficiency Syndrome (SCID) wherein T, B or both cell types may be depleted in humans. It is the existence of diseases affecting the immune system, such as AIDS and SCID, which has created the need for animal model systems to study the epitology and potential treatment of such disease states.

T lymphocytes recognize antigen in the context of self Major Histocompatibility Complex (MHC) molecules by means of the T cell receptor (TCR) expressed on their cell surface. The TCR is a disulfide linked heterodimer noncovalently associated with the CD3 complex (Allison, J.P., et al. (1987) Ann. Rev. Immunol. 5, 503). Most T cells carry TCRs consisting of α and β glycoproteins. T cells use mechanisms to generate diversity in their receptor molecules similar to those operating in B cells (Kronenberg, M., et al. (1986) Ann. Rev. Immunol. 4, 529; Tonegawa. S (1983) Nature 302, 575). Like the immunoglobulin (Ig) genes, the TCR genes are composed of segments which rearrange during T cell development. TCR and Ig polypeptides consist of amino terminal variable and carboxy terminal constant regions. The variable region is responsible for the specific recognition of antigen, whereas the C region functions in membrane anchoring and in transmitting of the signal that the receptor is occupied, from the outside to the inside of the cell. The variable region of the Ig heavy chain and the TCR β chain is encoded by three gene segments, the variable (V), diversity (D) and joining (J) segments. The Ig light chain and the TCR α chain contain variable regions encoded by V and J segments only.

The V, D and J segments are present in multiple copies in germline DNA. The diversity in the variable region is generated by random joining of one member of each segment family. Fusion of gene segments is accompanied by insertion of several nucleotides. This N-region insertion largely contributes to the diversity, particularly of the TCR variable regions (Davis and Bjorkman (1986) Nature 334, 395), but also implies that variable gene

segments are often not functionally rearranged. The rearrangement of gene segments generally occurs at both alleles. However, T and B cells express only one TCR or Ig respectively and two functionally rearranged genes within one cell have never been found. This phenomenon is known as allelic exclusion.

During B cell development the rearrangement process starts at both heavy chain gene alleles. First a D segment is fused to a J segment followed by ligation of a V segment to the DJ join. If the VDJ joining is productive, further rearrangement of the other heavy chain allele is blocked, whereas rearrangement of the light chain loci is induced (Reth, M., et al. (1985) Nature 317, 353).

In both B and T cells, partially (DJ) and completely (VDJ) rearranged genes reportedly are transcribed giving rise to two differently sized RNA molecules (Yancopoulos, G., et al. (1986) Ann. Rev. Immunol. 4, 339; Born, W., et al. (1987) TIG 3, 132). In B cells the DJ transcripts can be translated into a D μ -chain, a truncated form of the Ig μ heavy chain that lacks a V segment derived sequence. In general, the D μ -chain is present in minor amounts, if at all. However, in one subclone (P4-11) of the 300-19 cell line, a transformed pre-B cell line which differentiates in vitro to Ig producing B cells, the expression of the D μ -chain is reportedly very high (Reth, M., et al. (1985) Nature 317, 353). This reference also reports that the heavy chain gene alleles in the P4-11 clone are blocked at the DJ rearrangement stage in cell culture and that such cells show a very high frequency of light chain gene rearrangements. This has led to the suggestion that the D μ protein

contains some of the regulatory determinales necessary for gene assembly (Yancopoulos, G., et al. (1986) Ann. Rev. Immunol. 4, 339, 356).

Transgenic mice containing functionally rearranged
5 Ig genes reportedly have been used in studying several aspects of Ig gene expression, e.g. tissue specific expression, the mechanism of segment rearrangement, allelic exclusion and repertoire development (Storb, U. (1987) Ann. Rev. Immunol. 5,
10 151). It has also been reported that the transgenic heavy chain polypeptide only inhibits the complete VDJ rearrangement of endogenous heavy chain genes if it contains a transmembrane domain (Storb, 1987; Iglesias, A., et al. (1987) Nature 330, 482;
15 Nussenzweig, M., et al. (1987) Science 236, 816; Nussenzweig, M., et al. (1988) J. Exp. Med. 167, 1969).

Recently, the inventors reported that functionally rearranged TCR β genes can be appropriately expressed
20 in transgenic mice (Krimpenfort, P., et al. (1988) EMBO J. 7, 745). This functional TCR β chain gene prevents expression of endogenous β genes by inhibiting complete VDJ joining (Uematsu, Y., et al. (1988) Cell 52, 831).

25 Two different types of T cells are involved in antigen recognition within the self MHC context. Mature T helper cells (CD4⁺CD8⁻) recognize antigen in the context of class II MHC molecules, whereas cytotoxic T cells (CD4⁻CD8⁺) recognize antigen in the
30 context of class IMHC determinants (Swain, S.L. (1983) Immun. Rev. 74, 129-142; Dialynas, P.D., et al. (1983) Immun. Rev. 74, 29-56). It has been reported that class II- specific CD4⁺CD8⁻ helper T

-6-

cells (also referred to as T4 cells) fail to develop in mice neonatally treated with anti-class II MHC monoclonal antibody (Kruisbeek, A.M., et al. (1983) J. Exp. Med. 157, 1932-1946; Kruisbeek, A.M., et al. (1985) J. Exp. Med. 161, 1029-1047). Similarly, it has recently been reported that mice chronically treated with anti-class I MHC monoclonal antibody from birth have a significantly reduced population of CD4⁻CD8⁺ cells and cytotoxic T cell precursors (Marusic-Galesic, S., et al. (1988) Nature 333, 180-183). Although selected T cell populations apparently can be produced by such methods, continuous administration of antibody is required which often results in adverse side effects in such mice.

A different strategy to deplete specific cell lines has recently been identified wherein specific cell destruction is induced by administration of a toxic metabolite. Specifically, transgenic mice reportedly were produced containing a Herpes Simplex Virus Thymidine Kinase (HSV-TK) transgene fused to the Ig promoter/enhancer. Transgenic cells that express the HSV-TK are not affected. However, upon administration of a nucleoside analog that can be phosphorylated by the transgenic HSV-TK gene, replicating cells expressing the HSV-TK gene are killed (Heyman, et al. (1988) UCLA Symposia on Molecular and Cellular Biology, 73, 199).

Another approach to depletion of specific cell types has been reported using tissue specific expression of a bacterial toxin. Specifically, mice carrying an elastase/diphtheria toxin A (DT-A) fusion gene lacked a normal pancreas (Palmer, et al. (1987) Cell 50, 435). In addition, it has been reported that

microphthalmia in transgenic mice resulted from the introduction of the DT-A gene fused to the $\alpha 2$ -crystallin promoter (Bretman, et al. (1987) Science 238, 1563).

5 Transgenic mice reportedly have also been constructed that express an $\alpha\beta$ TCR in a large fraction in their T cells which is specific for a minor histocompatibility antigen (H-Y) present on male, but not female, cells (Kisielow, P., et al. (1988) Nature
10 333, 742-746). This very recent reference reports that cells containing the TCR for the H-Y antigen were frequent in female but not in male transgenic offspring. The $\alpha\beta$ TCR in such transgenic mice
15 apparently contains all the segments and regions required for a functional TCR.

The references discussed above are provided solely for the disclosure prior to the filing date of the present application and nothing herein is to be construed as an admission that the inventors are not
20 entitled to antedate such disclosures by virtue of prior invention.

Given the state of the art, it is apparent that a need exists for animal model systems to study diseases which effect the immune system including
25 infectious diseases such as AIDS. Accordingly, it is an object herein to provide transgenic non-human animals and methods for making the same which have a phenotype characterized by the substantial depletion of a mature lymphocytic cell type otherwise naturally
30 occurring in the species from which the transgenic is derived.

It is also an object herein to provide transgenic non-human animals substantially depleted in mature T cells or plasma cells.

5 It is a further object herein to provide transgenic mice substantially depleted in mature T cells or plasma cells.

Still further, it is an object herein to provide transgenes capable of producing such transgenic non-human animals.

10 Further, it is an object herein to provide methods for producing transgenic non-human animal having at least one of the above identified phenotypes.

Summary of the Invention

15 The invention is based on the discovery that transgenic non-human animals depleted in a lymphatic cell type can be produced by disrupting the expression of a functional lymphocytic polypeptide required for maturation of the lymphocytic cell type. This lymphocytic polypeptide is otherwise expressed
20 by the non-human animal from which the transgenic animal is derived.

In one aspect, the invention provides transgenic non-human animals having a phenotype characterized by the substantial depletion of a mature lymphocytic
25 cell type otherwise naturally occurring in the species from which the transgenic animal is derived. This phenotype is conferred in the transgenic animal by a transgene contained in at least the precursor stem cell of the lymphocytic cell type which is
30 depleted. The transgene comprises a DNA sequence encoding a lymphatic polypeptide variant which

inhibits formation of the depleted lymphocytic cell type. Generally, such inhibition occurs when the lymphatic polypeptide variant is expressed in a precursor to the lymphocytic cell type.

- 5 In those cases where the lymphatic polypeptide variant is expressed, the variant is believed to be capable of suppressing expression of at least one set of cognate endogenous alleles normally expressed during differentiation of the precursor stem cell to
- 10 the mature lymphocytic cell type. The lymphatic polypeptide variant, however, lacks a functional domain necessary for maturation of the lymphocytic cell type which would otherwise be provided by either or both of the suppressed endogenous alleles.
- 15 Within the context of transgenic animals deficient in T cells, the transgene encodes a lymphatic polypeptide variant comprising a portion of a TCR β chain. The transgene encoding the TCR β variant chain typically retains sequences encoding at least the
- 20 transmembrane sequence found in the C region of a naturally occurring TCR β chain. This sequence may be operably linked to an appropriate signal sequence. It lacks, however, sequences encoding all or part of the variable region. The C region contained by such
- 25 a lymphatic polypeptide variant is capable of suppressing the expression of endogenous TCR alleles thereby preventing the membrane expression of functional heterodimeric TCRs. Normal T cell maturation is thereby abrogated.
- 30 In the case of non-human transgenic animals substantially depleted in antibody secreting plasma cells, the transgene similarly encodes a lymphatic polypeptide variant containing at least the

transmembrane sequence of the C region of the Ig heavy chain. A signal sequence may also be operably linked to the transgene encoding the lymphatic polypeptide variant.

- 5 The invention also includes transgenes comprising a DNA sequence encoding a lymphatic polypeptide variant.

Further, the invention includes a method for producing a transgenic non-human animal substantially
10 depleted in a mature lymphocytic cell type. The method comprises introducing a transgene into an embryonal target cell. The transgene encodes a lymphatic polypeptide variant and is capable of inhibiting formation of a mature lymphocytic cell
15 type. The thus transformed transgenic embryonal target cell is thereafter transplanted into a recipient female parent from which offspring having the desired phenotype are identified.

Detailed Description of the Drawings

- 20 Fig. 1A is a schematic representation of the $\alpha\beta$ T cell receptor.

Fig. 1B depicts the $\alpha\beta$ T cell receptor variant containing a deletion of the variable region of the β chain.

- 25 Fig. 2 depicts the genomic organization of the T cell receptor loci in mouse.

Fig. 3A and 3B show the structure of the T cell receptor β transgene including nucleotide sequence of the variable region and restriction map of clone cos
30 HY9-1.14-5.

-11-

Fig. 4 is a Southern blot analysis of offspring mice derived from a cross of the female transgenic mouse 93 with C57L males.

5 Fig. 5 demonstrates the surface expression of F23.1-positive β chains on T cells from transgenic and normal mice.

Fig. 6 demonstrates the tissue specificity of transgene transcription.

10 Fig. 7 is a gel electrophoretic analysis of immunoprecipitated β chains from transgenic T lymphocytes before and after preclearing with F23.1 monoclonal antibodies.

15 Fig. 8 demonstrates the modulation of surface expression of CD3 molecules on various T cells by V β 8-specific F23.1 monoclonal antibodies.

Fig. 9 demonstrates the inhibition of cytolytic activity of transgenic T cells by V β 8-specific F23.1 monoclonal antibodies.

20 Fig. 10 demonstrates that endogenous β genes in transgenic mouse 93.2 show predominantly partial D β 1-J β rearrangements.

Figs. 11 and 12 depict the construction of Δ V-TCR β and Δ V α -TCR β transgenes.

25 Fig. 13 demonstrates the expression of the Δ V α -TCR β transgene.

-12-

Fig. 14 is a surface marker staining profile of splenocytes from transgenic mouse #1670 and a control mouse.

Fig. 15 depicts the amino acid and known DNA sequence for transgenes TCR β and Δ V-TCR β based on partial cDNA and genomic sequence data.

Detailed Description of the Disclosure

The "non-human animals" of the invention comprise any non-human animal having an immune system capable of producing a humoral and/or cell-mediated immune response. Such non-human animals include vertebrates such as rodents, non-human primates, sheep, dog, cow, amphibians, reptiles, etc. Preferred non-human animals are selected from the rodent family including rat and mouse, most preferably mouse.

The "transgenic non-human animals" of the invention are produced by introducing "transgenes" into the germline of the non-human animal. Embryonal target cells at various developmental stages can be used to introduce transgenes. Different methods are used depending on the stage of development of the embryonal target cell. The zygote is the best target for micro-injection. In the mouse, the male pronucleus reaches the size of approximately 20 micrometers in diameter which allows reproducible injection of 1-2pl of DNA solution. The use of zygotes as a target for gene transfer has a major advantage in that in most cases the injected DNA will be incorporated into the host genome before the first cleavage (Brinster, et al. (1985) Proc. Natl. Acad. Sci. USA 82, 4438-4442). As a consequence, all cells of the transgenic non-human animal will carry the

-13-

incorporated transgene. This will in general also be reflected in the efficient transmission of the transgene to offspring of the founder since 50% of the germ cells will harbor the transgene. Micro-
5 injection of zygotes is the preferred method for incorporating transgenes in practicing the invention.

Retroviral infection can also be used to introduce transgene into a non-human animal. The developing non-human embryo can be cultured in vitro to the
10 blastocyst stage. During this time, the blastomeres can be targets for retroviral infection (Jaenich, R. (1976) Proc. Natl. Acad. Sci USA 73, 1260-1264). Efficient infection of the blastomeres is obtained by enzymatic treatment to remove the zona pellucida
15 (Hogan, et al. (1986) in Manipulating the Mouse Embryo, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.). The viral vector system used to introduce the transgene is typically a replication-defective retrovirus carrying the
20 transgene (Jahner, et al. (1985) Proc. Natl. Acad. Sci. USA 82, 6927-6931; Van der Putten, et al. (1985) Proc. Natl. Acad. Sci USA 82, 6148-6152). Transfection is easily and efficiently obtained by culturing the blastomeres on a monolayer of virus-
25 producing cells (Van der Putten, supra; Stewart, et al. (1987) EMBO J. 6, 383-388). Alternatively, infection can be performed at a later stage. Virus or virus-producing cells can be injected into the blastocoele (Jahner, D., et al. (1982) Nature 298,
30 623-628). Most of the founders will be mosaic for the transgene since incorporation occurs only in a subset of the cells which formed the transgenic non-human animal. Further, the founder may contain various retroviral insertions of the transgene at
35 different positions in the genome which generally

will segregate in the offspring. In addition, it is also possible to introduce transgenes into the germ line, albeit with low efficiency, by intrauterine retroviral infection of the midgestation embryo
5 (Jahner, D. et al. (1982) supra).

A third type of target cell for transgene introduction is the embryonal stem cell (ES). ES cells are obtained from pre-implantation embryos cultured in vitro and fused with embryos (Evans,
10 M.J., et al. (1981) Nature 292, 154-156; Bradley, M.O., et al. (1984) Nature 309, 255-258; Gossler, et al. (1986) Proc. Natl. Acad. Sci USA 83, 9065-9069; and Robertson, et al. (1986) Nature 322, 445-448). Transgenes can be efficiently introduced into the ES
15 cells by DNA transfection or by retrovirus-mediated transduction. Such transformed ES cells can thereafter be combined with blastocysts from a non-human animal. The ES cells thereafter colonize the embryo and contribute to the germ line of the
20 resulting chimeric animal. For review see Jaenisch, R. (1988) Science 240, 1468-1474.

As used herein, a "transgene" is a DNA sequence introduced into the germline of a non-human animal by way of human intervention such as by way of the
25 above described methods. The transgenes of the invention include DNA sequences which are capable of suppressing cognate endogenous alleles. Further, such transgenes are capable of either facilitating or inhibiting the maturation of a lymphatic cell type.
30 Such transgenes comprise DNA sequences encoding either a "lymphatic polypeptide" or "lymphatic polypeptide variant" which may be expressed in a transgenic non-human animal.

A lymphatic polypeptide corresponds to a naturally occurring polypeptide expressed selectively by lymphatic tissue and includes the wide variety of polypeptide chains associated with immunoglobulin production by plasma cells and TCRs produced by mature T cells. Such lymphatic polypeptides are produced by mature lymphocytic cell types after differentiation. Such differentiation involves the functional rearrangement of numerous gene regions and segments to form the DNA sequence encoding the naturally occurring lymphatic polypeptide (described in more detail hereinafter).

The term "cognate endogenous alleles" refers to alleles in the genome of the transgenic non-human animal which are closely related to the transgene introduced therein. Thus, for example, the alleles involved in TCR β chain production in mouse T cells are cognate endogenous alleles which may be suppressed by a transgene comprising a DNA sequence encoding a TCR β chain from mouse or other species. Further, such cognate endogenous alleles may include the $\gamma\delta$ alleles involved in $\gamma\delta$ TCR production.

The transgene encoding a lymphatic polypeptide, in addition to suppressing cognate endogenous alleles, is also capable of facilitating the maturation of the particular lymphocytic cell type which would normally express the cognate endogenous alleles if the lymphatic polypeptide is expressed. Thus, transgenic mice containing a transgene encoding a functionally rearranged β TCR chain contain mature T lymphocytes which express $\alpha\beta$ TCRs containing β chains encoded by the TCR β transgene.

Transgenes encoding lymphatic polypeptide variants, however, suppress cognate endogenous alleles and inhibit the maturation of the lymphocytic cell type which normally expresses such cognate endogenous alleles. The parameters which define the lymphatic polypeptide variant may be defined structurally and functionally.

As indicated in the Examples, a functionally rearranged mouse TCR fusion β gene which is expressed in a transgenic mouse, suppresses the expression of the cognate endogenous alleles encoding the β chain TCR gene. See also Uematsu, Y., et al. (1988) Cell 52, 831-841. The suppression of such expression apparently results from the blocking of the rearrangement process of the endogenous β genes during differentiation to the mature T lymphocyte. Partial D-J rearrangements are found in such transgenic animals while complete VDJ rearrangements are not detected. Such transgenic mice, however, do produce functional mature T cells. The $\alpha\beta$ TCRs on such transgenic mice are homogeneous with respect to the β chain encoded by the transgene. They are, however, fully functional, at least in a variety of allogenic responses, presumably due to the diversity produced in such receptors by way of endogenous α chain rearrangements.

When, however, transgenic mice are produced containing a functionally rearranged TCR β gene which has had approximately 90% of the variable region deleted (Δ TCR β), such mice failed to form a functional thymus and substantial T cell depletion was observed. Thus, disruption of the expression of functional TCR β chains is sufficient to cause T cell depletion. The mechanism for this T cell depletion

is not known. However, it is assumed that the TCR β gene encodes a lymphatic polypeptide having at least two functional domains related to the mechanism of T cell depletion when transgenes containing variable region deletions are used. One domain functions to suppress the expression of cognate endogenous alleles (such as the TCR β chain alleles) presumably by a mechanism analogous to allelic exclusion observed during normal differentiation. The second functional domain is required for T cell maturation. In the case of the complete TCR β gene both functional domains are present in the transgenic TCR β chain as evidenced by the suppression of endogenous β alleles and the normal maturation of T lymphocytes. However, in the case of the transgene ΔV -TCR β (containing a deletion of approximately 90% of the variable region), the transgenic ΔV -TCR β polypeptide contains only a first functional domain capable of suppressing endogenous β chain alleles. A sufficient portion of the second functional domain is deleted such that T cell maturation is inhibited. See Fig. 1B.

Thus, structurally, the transgene, in one aspect of the invention, encodes a lymphatic polypeptide variant comprising a lymphatic polypeptide wherein all or part of the variable region is deleted. Preferably, at least part of the V segment of the V region is deleted. However, the deleted sequences may also include part of the D and/or J segment of the variable region of the lymphatic polypeptide.

The transgene encoding the lymphatic polypeptide variant also can be defined functionally. This definition is based on the two functional domains present in the naturally occurring lymphatic polypeptide. By analogy to the TCR β chain, the first

-18-

functional domain is capable of suppressing the expression of cognate endogenous alleles whereas the second functional domain is capable of inducing the maturation of a particular lymphocytic cell type.

5 The lymphatic polypeptide variant contains the first functional domain but lacks the second. Although defined functionally, such domains can be structurally ascertained by way of the methodology disclosed in Examples 1 and 2 herein. For example,

10 the first functional domain involved in the suppression of cognate endogenous alleles may be ascertained by generating a family of transgenes encoding specific deletions of portions of the lymphatic polypeptide. Transgenic mouse lines,

15 containing each of these transgenes, for example, may then be analyzed to ascertain the suppression of endogenous alleles as described in Example 1. Those transgenes containing deletions which do not induce suppression of the endogenous alleles define the

20 functional domain responsible for suppression of endogenous expression.

Once such structural domains are defined, the second functional domain may be defined structurally by generating a series of transgene constructs

25 containing deletions of various portions of the lymphatic polypeptide not defining the first functional domain. Thus, sequential deletions (i.e., deletions of about 10-30 base pairs through, for example, the variable region) or progressive

30 deletions (i.e., a set of deletions comprising deletion of approximately 10-20 base pairs to, for example, all of the DNA sequence encoding the variable region) can be used in conjunction with the methods of Example 2 to define the second functional

35 domain required for lymphocyte maturation. Thus,

-19-

lymphatic polypeptide variants and the transgenes encoding the same can be readily defined by such methods.

As indicated, the lymphatic polypeptide variant
5 encoded by the transgene in one embodiment of the invention should be expressed in a precursor of the mature lymphocyte to be depleted in the transgenic animal. To be expressed, the transgene DNA sequence must contain regulatory sequences necessary for
10 transcription of the transgene in the precursor cell type in the transgenic animal. Further, various regulatory sequences necessary for translation and if necessary processing of the mRNA transcripts encoded by the transgene are required. Such regulatory
15 sequences include promoter sequences to control transcription, ribosome binding sites to facilitate translation, splice acceptor and splice donor sequences for intron excision, polyadenylation sequences, transcription and translational
20 termination signal and enhancer sequences which facilitate the efficient expression of the lymphatic polypeptide variant. In most cases, a secretory signal sequence will also be included to facilitate the association of the lymphatic polypeptide variant
25 with the membrane of the cell expresssing the transgenic polypeptide. Transgenes therefore encode all of the regulatory sequences necessary to express the lymphatic polypeptide or lymphatic polypeptide variant encoded by the transgene. Of course, each of
30 the regulatory sequences and encoding sequences are operably linked to produce a functional transgene capable of expressing the transgenic polypeptides in the non-human transgenic animal.

"Operably linked" when describing the relationship between two DNA or polypeptide sequences simply means that they are functionally related to each other. For example, a signal or leader sequence is operably
5 linked to a peptide if it functions as a signal sequence participating in the insertion of the immature form of the protein into a cell membrane. Similarly, a promoter is operably linked to a coding sequence if it controls the transcription of the
10 sequence; a ribosome binding site as operably linked to a coding sequence if it is positioned so as to permit translation, etc.

Transgenes are derived, for example, from DNA sequences encoding at least one polypeptide chain of
15 a T cell receptor (TCR) or one polypeptide chain of the immunoglobulin (Ig) molecule. Preferably, a modified form of the β or γ chain of the TCR and most preferably the β chain of the TCR is used as a transgene to inhibit the formation of mature T cells.
20 In the case of B cells, a derivative of the heavy chain of the Ig molecule is preferred to inhibit the formation of antibody producing plasma cells derived from B cells. Generally, such transgenes are derived by deleting from the DNA sequence encoding a
25 functionally rearranged β chain, γ chain or heavy chain polypeptide, all or part of the DNA sequence encoding the variable region of such molecules. Preferably all of the variable region is deleted although small segments of 5' sequences encoding an
30 N-terminal portion of the V segment and 3' sequences encoding the C-terminal portion of the J segment may be retained in the transgene. At the very least, all or part of the variable segment should be deleted. Thus, transgenes generally comprise C regions of the
35 β chain, γ chain or heavy chain polypeptides of TCR

and Ig molecules respectively but may also include additional sequences encoding all or part of the J and D segments of the variable region.

Transgenic mice containing a transgene encoding a
5 TCR β chain having its variable regions deleted do
not contain detectable T cells. Further, transgenic
mice containing immunoglobulin heavy chain wherein
all or part of the variable region is deleted are
expected to be incapable of producing plasma cells
10 which secrete immunoglobulins. This is because
during B cell development, B cells rearrange their Ig
heavy chain genes first. Once a functionally
rearranged Ig heavy chain gene is generated, light
chain rearrangement starts. This eventually will
15 result in the production of complete IgM molecules
containing two heavy and two light chains. Such IgMs
are not secreted since their C regions contain a
membrane anchoring domain. During further
development, the B cells switch the use of the
20 constant region to other constant regions that do not
encode a transmembrane domain, e.g., IgG. When this
switch occurs, they become plasma cells which secrete
large amounts of specific immunoglobulin. In order
to develop into plasma cells, the IgM producing B
25 cells must interact with other cells of the immune
system via the IgM located on the B cell surface.
Since the heavy chain of the IgM variant contains the
deletion of all or part of the variable region,
transgenic non-human animals containing a transgene
30 encoding such a deleted heavy chain should not be
able to produce B cells which can interact with the
immune system to form the mature plasma cell type.

Alternatively, the transgene may comprise a DNA
sequence encoding a lymphatic polypeptide variant

which is incapable of being expressed in the precursor stem cell or latter precursor of the lymphocytic cell type to be depleted. Such DNA sequences may contain one or more mutations involving the insertion or deletion of one or more nucleotides that may, for example, result in a frame shift or nonsense mutation (e.g., stop codon) that prevents all or part of the expression of the transgene. The transgene, however, has sufficient sequence homology with a cognate endogenous allele such that when introduced into an ES target cell it may homologously recombine with such an allele in the ES cell to disrupt its expression. See, e.g., Kirk and Capecchi (1978) Cell 51, 503-512. After identification and selection of ES cells containing the transgene in the targeted allele, a transgenic non-human animal may be produced by colonizing an appropriate embryo with the selected ES cell.

In the case of T cells, the $C\beta 1$ and $C\beta 2$ alleles for the $TCR\beta$ chain and most preferably the $C\alpha$ allele for the $TCR\alpha$ chain are targeted for insertion of a transgene which disrupts expression of the allele. For example, in the case of the $C\alpha$ allele, once a genotype is identified containing a transgene disrupting $C\alpha$ expression, cross-breeding can be used to produce transgenic animals homozygous for the $C\alpha^-$ genotype. Such homozygotes should also be deficient in mature T cells since they are incapable of producing the functional $\alpha\beta TCR$ required for T cell maturation.

A similar approach can be used to produce plasma cell deficient transgenic animals. In that case the transgene is targeted to disrupt the expression of $C\mu$ and $C\delta$ portions of the IG heavy chain.

-23-

Transgenic non-human animals depleted in one or more mature lymphocytic cell types, such as transgenic mice depleted in mature T cells or plasma cells, have multiple uses. For example, T cell depleted mice do not have a cell-mediated immune response. They are therefore suitable for testing drugs that interfere with cell-mediated immunity for side effects and complications, e.g., kidney damage, etc. Thus, antibiotics, anti-viral drugs, antifungal agents and immunosuppressive drugs such as cyclosporine may be administered to such mice to ascertain their effects.

In addition, such T cell deficient mice do not have T4 (helper) cells used in the B cell humoral response. They can, therefore, be used for testing the effects of passive immunization. Further, the lack of a cell-mediated immune response renders such transgenic mice prone to develop tumors since they are not protected by immuno-surveillance. They therefore can be used to test the carcinogenicity of various materials. In addition, materials may be tested in such animals which may confer protection against the development of neoplasms.

Further, such T cell depleted mice can be used as model systems for AIDS and SCID involving the depletion of T cells.

Plasma cell deficient transgenic mice, in combination with T cell depleted mice, offer a good model system to study the effect of the depletion of the humoral immune response, cell-mediated immune response or both. Further, such plasma cell depleted and/or T cell depleted mice can be used as a model system to

-24-

study SCID syndromes involving the depletion of B, T and/or B and T cell types.

The genes encoding the various segments and regions which may be used in the invention are well characterized. The TCRs associated with T cells represent an enormous percent of clonally varying molecules with the same basic structure. The TCR is a heterodimer of 90kd consisting of two transmembrane polypeptides of 45kd each connected by disulfide bridges (Fig. 1A) (Samuelson, et al. (1983) Proc. Natl. Acad. Sci. USA 80, 6972; Acuto, et al. (1983) Cell 34, 717; MacIntyre, et al. (1983) Cell 34, 737). For most T cells, the two polypeptides are referred to as the α and β chain. Like the heavy and light chains of the immunoglobulins, the α and β chains have variable (V) and constant (C) regions (Acuto, et al. (1983) supra; Kappler, et al. (1983) Cell 35, 295). The V region is responsible for antigen recognition and the C region is involved in membrane anchoring and signal transmission. A small percentage of lymphoid T cells, however, have a different set of TCRs comprising different polypeptide chains referred to as γ and δ chains (Borst, et al. (1987) Nature 325, 683; Brenner, et al. (1987) Nature 325, 689; Bank, et al. (1986) Nature 322, 179; Pardoll, et al. (1987) Nature 326, 79).

Using subtractive hybridization procedures, cDNA clones encoding the TCR polypeptide chains have been isolated (Hendrick, et al. (1984) Nature 308, 149; Hendrick, et al. (1984) Nature 308, 153; Yanagi, et al. (1984) Nature 308, 145; Saito, et al. (1987) Nature 325, 125; Chien, et al. (1984) Nature 312, 314). Sequence analysis of these cDNA clones reveal

-25-

the complete primary sequence of the TCR polypeptides. The TCR polypeptides are all similar to each other and to the immunoglobulin polypeptides. For review see Davis and Bjorkman
5 (1988) supra.; and Kronenberg, et al. (1986) Ann. Rev. Immunol. 4, 529). The variable region of the TCR chains is composed of a variable region (V) and a constant region (C). The variable region of the TCR chains is further subdivided into a variable (V) and
10 joining (J) segments. In addition, the variable region of the β and δ chains also contains a diversity (D) segment interposed between the V and J segments. At similar positions as in the immunoglobulins, hypervariable regions are present in
15 the variable region of the TCR. Two hypervariable regions are encoded in the V region whereas one is represented by the junction between the V and J or V, D and J segments. The constant region of the TCR chains is composed of four functional regions often
20 encoded by different exons (Davis and Bjorkoran (1988) supra.). The TCR C regions show typical immunoglobulin-like characteristics such as the presence of cystine residues involved in the linking of two β sheets which probably fold into
25 immunoglobulin-like domains. All TCR chains have a hydrophobic transmembrane region in which a highly conserved lysine residue is present.

The availability of TCR cDNAs permits an analysis of the genomic organization of the murine and human TCR
30 genes. The TCR genes show a segmental organization similar to the immunoglobulin genes. Fig. 2 shows a schematic representation of the four TCR gene loci. In the β chain gene locus, two nearly identical C β regions are tandemly arranged, each preceded by one
35 D and six J segments (Kronenberg, et al. (1986),

supra; Davis, M. (1985) Rev. Immunol. 3, 537). The β locus also contains approximately 20 to 30 V segments (Barth, et al. (1985) Nature 316, 517; Behlkey, et al. (1985) Science 229, 566), one of which is located 3' to the C regions in opposite orientation (Malissen, et al. (1986) Nature 319, 28). The γ locus is less diverse, containing three J_γ - C_γ regions and only a limited number of V segments (Hayday, et al. (1985) Cell 40, 259; Chien, et al. (1987) Nature 330, 722). The α and δ loci overlap each other in that many of the δ coding segments are located within the segments of the α gene (Chien, et al., supra; Elliott, et al. (1988) Nature 331, 627). As a consequence, the δ segments are deleted in most $\alpha\beta$ bearing T cells (Lindsten, et al. (1987) J. Exp. Med. 166, 761). Thus, the α and γ chain genes are encoded by variable and joining segment and constant regions whereas the β and δ chain genes contain variable, diversity and joining segment and a constant region. During somatic development of the T cell, a functional TCR gene is formed by rearrangement of these segments and regions. This process is the basis for T cell receptor diversity. The following strategies have been postulated for T cell diversification: (1) multiple germline V and J segments (Kronenberg, et al. (1986) supra; (2) D segments that can be translated in all three reading frames in the case of β and δ genes (Goverman, et al. (1985) Cell 40, 859; Elliott, et al. (1988), Nature 331, 627); (3) combinatorial joining of V, D and J segments; (4) the random addition of nucleotides between the V, D and J segments (N-region addition) (Davis and Bjorkman (1988) supra); (5) the flexibility of several V segments with respect to the position of their 3' joining points (Davis and

Bjorkman, supra); and (6) the combinatorial joining of TCR chains.

As shown schematically in Fig. 2, the encoding segments for the TCR genes are scattered over large
5 arrays of chromosomal DNA. Like the immunoglobulin genes, specific V, D and J segments are fused together to generate a complete V coding region next to a C region. In B and T cells, the rearrangements are mediated by similar sequences flanking the
10 segments to be fused (Akira (1987) Science 238, 1134; Yancopoulos, et al. (1986) supra.). These sequences consist of conserved heptamer and monomer stretches spaced by 12 or 23 nucleotides. Depending on the orientation of the segments being joining, looping
15 out/deletion or inversion of large genomic DNA can occur (Fujimoto (1987) Nature 327, 242; Okazaki, et al. (1987) Cell 49, 477; Mallisen, B., et al. (1986) supra). B and T cells probably use the same machinery for the assembly of Ig and TCR since B
20 cells rearrange transfected TCR segments in the same way as transfected Ig gene segments (Yancopoulos, et al. (1986) supra). The TCR β , γ and δ genes are rearranged and transcribed first, followed by the TCR α gene (Chien, et al. (1987) supra; Pardoll, et al. (1987) Nature 326, 79; Raulet, et al. (1985) Nature
25 312, 36; Samelson, et al. (1985) Nature 315, 765; Snodgrass, et al. (1985) Nature 315, 232).

The regulation of immunoglobulin gene assembly and expression is intrinsically related to the progression
30 of B-cell precursors to the B cell differentiation stage. For a review of the mechanism of Ig variable region gene assembly and regulatory mechanisms to control genomic rearrangements, see Yancopoulos, et al. (1986) supra. Ig gene assembly in B cells

closely parallels that for TCR assembly in T cells. In addition, a number of transgenic mice containing a number of immunoglobulin genes have been prepared to further study the mechanism of Ig expression, allelic
5 exclusion and rearrangement mechanisms (Storb (1987) Ann. Rev. Immunol. 5, 151-174).

Thus, the $\alpha\beta$ and $\gamma\delta$ TCRs of T cells and the heavy and light chains of Ig molecules in B cells have been well characterized. As indicated, the transgenes of
10 the present invention are derived from such DNA sequences, preferably those encoding the β and γ chains of the T cell receptors, most preferably the β chain, and the heavy (H) chain of the Ig molecule from B cells. Such DNA may be obtained from the
15 genome of somatic cells and cloned by well established technology. Such cloned DNA sequences may thereafter be further manipulated by recombinant techniques to construct the transgenes of the present invention. However, a more efficient and preferred
20 methodology is to clone a functionally rearranged β or γ TCR gene or a functionally rearranged heavy Ig gene. Such cloned functionally rearranged genes generally will have all necessary regulatory and secretory sequences for expression of the
25 functionally rearranged gene in the B or T lymphocytes of the animal from which the sequence is derived, including introns which are necessary for the efficient expression of transgenes (Brinster, et al. (1988) Proc. Natl. Acad. Sci. USA 85, 836). Such
30 DNA sequences can be derived by generating cultures of T or B cells from mature animals or animals which have been challenged with an antigen. The DNA from one or more such clones may be used to generate a genomic DNA library in hosts such as λ gt11 and EMBL 3
35 following published procedures (Young, R.A., et al.

(1983) Proc. Natl. Acad. Sci. USA 80, 1194-1198; Frischauf, et al. (1983) J. Mol. Biol. 170, 827-842). Such libraries may then be screened with appropriate probes to particular regions or segments of the β and/or γ genes for TCRs or the heavy gene for Ig molecules depending on the source of the genomic DNA. Such probes preferable are specific for a part of one of the C regions of the molecule of interest. cDNA clones from the same cellular clone may also be generated and identified by similar probes. The sequences of such cDNA clones facilitates the identification of the coding sequence of the genomic clone thereby permitting the identification of putative introns and regulatory sequences.

15 If a particular genomic clone does not contain the entire coding region for the functionally rearranged chain or if it does not contain regulatory or enhancer regions typically located in 5' or 3' flanking regions, that DNA sequence may be recombined with other clones to provide such sequences. Thus, for example, a genomic clone from the T cell clone B6.2.16 (described in more detail in the Examples) has flanking sequences comprising 2kb 5' and 2kb 3' from the coding region for the mouse TCR β gene.

20 This DNA sequence is inactive in transgenic mice because it lacks a 550 base pair enhancer sequence located 5kb downstream from the C β 2 locus (Krimpenfort, P., et al. (1988) EMBO J. 7, 745-750). A fusion construct was therefore prepared containing a larger extension of the 3' end. A cosmid clone derived from BALB7c liver DNA containing the C β 2 and downstream region was used to generate this fusion construct which is active in transgenic mice (Krimpenfort, P., et al. (1988) supra). Similar methods may be used to generate fusion genes

containing all the necessary DNA sequences for regulation of the expression of the DNA sequences encoding the particular polypeptide chain of interest.

5 The DNA sequences from which the transgene is derived are preferably obtained from the functionally rearranged genome of the same species of animal into which the transgene will be introduced. Thus, functionally rearranged β and γ genes from mouse TCR
10 are preferred for making transgenes for use in transgenic mice. The invention, however, is not limited to transgenes derived from the same species of animal used to form transgenic non-human animals. It has recently been shown that two independent lines
15 of transgenic mice containing either the human heavy chain of HLA class I antigen or the light (β_2 -microglobulin) chain of human HLA class I antigen can be crossed to produce a functional HLA class I antigen which is biochemically indistinguishable from
20 the same antigen expressed on human cells (Krimpenfort, P., et al. (1987) EMBO J. 6, 1673-1676. Other examples of heterologous transgenic expression include the heterologous regulatory and structural sequences disclosed in EPO Publication No. 0247494
25 and PCT Publication No. WO88/00239. Thus, DNA sequences from species different from that of the transgenic animal ("heterologous DNA") may be used to form the transgenic non-human animals of the present invention. Such heterologous sequences include
30 regulatory and secretory sequences as well as structural DNA sequences which when modified encode heterologous lymphatic polypeptide variants capable of inhibiting the formation of a mature lymphocytic cell type. The only limitation on the use of such
35 heterologous sequences is functional. The

heterologous regulatory sequences must be utilized by the transgenic animal to efficiently express sufficient amounts of the lymphatic polypeptide variant such that it is able to inhibit the formation of a mature lymphocytic cell type. Further, the heterologous lymphatic polypeptide variant when properly expressed in the transgenic animal must be capable of producing the desired depletion of a lymphocytic cell type. Thus, by analogy to expression of human HLA antigens on the cell surface of transgenic mice, a functionally rearranged human TCR β gene containing a variable region deletion can be used to express a human β TCR variant capable of suppressing the formation of mature T cells in transgenic mice. Further, it should be possible to mix homologous and heterologous DNA sequences (e.g., homologous regulators with heterologous structural genes and vice versa) to produce functional transgenes which may be used to practice the invention. Alternatively, heterologous DNA sequences encoding lymphatic polypeptide variant must be capable of inhibiting the expression of functional lymphatic polypeptide required for the maturation of a lymphocytic cell type by disrupting the expression of cognate endogenous alleles.

The following is presented by way of example and is not to be construed as a limitation on the scope of the invention. Further, all references referred to herein are expressly incorporated by reference.

30

Example 1SUPPRESSION OF COGNATE
ENDOGENOUS ALLELES OF TCR β

This example describes the construction of a transgene which when expressed is capable of

-32-

suppressing the expression of cognate endogenous alleles for the TCR β chain. When introduced into a mouse, the transgenic mouse so produced expresses TCR receptors substantially containing only the TCR β chain encoded by the transgene.

Establishment of the
Male-Specific B6.2.16 Clone

C57BL/6 female mice were immunized with 10^7 male spleen cells injected intraperitoneally. After 14 days, 2×10^7 female spleen cells were cultured with 2×10^7 x-irradiated male cells in 8ml of culture media. After 12 days, 10^6 viable female responder cells were restimulated with 10^7 x-irradiated male cells in 1ml of medium containing interleukin-2 (IL-2). Cells were cloned by culturing 0.3 cells with stimulators as described below. Clones were tested for cytolytic activity as described below, and clone B6.2.16 was selected after surface staining of various clones with the F23.1 antibody (von Boehmer and Haas, (1986) Meth. Enzymol. 132, 467-477; Staer, et al. (1985) J. Immunol. 134, 3994-4000) which binds to all three members of the V β 8 family (Behlke, et al. (1982) J. Exp. Med. 165, 257-267).

Isolation and Characterization of
 β Chain cDNA and Genomic Clones

Total RNA and high molecular weight DNA were isolated from cytotoxic T cell clone B6.2.16 and used to construct cDNA and genomic DNA libraries in λ gt11 and EMBL3, respectively, following published procedures (Young and Davis (1983) Proc. Natl. Acad. Sci. USA 80, 1194-1198; Frischauf, et al. (1983) J. Mol. Biol. 170, 827-842). Both libraries were screened with a C β probe corresponding to the sequence of the C β 2 region reported by Mallisen, et al. (1984) Cell 37,

1101. Positive clones were characterized by mapping with restriction endonucleases and hybridization with probes specific for the D β 1 gene segment, 1 1.4kp Pst1 fragment extending 1.25kb 5' to the D β 1 gene segment (Siu, et al. (1984) Nature 311, 344-350), and J β gene segment. A full-length cDNA clone and genomic clones, λ 8 for the functional allele and λ 3 for the nonfunctional allele, were subcloned into M13mpl8 for DNA sequence determination. Sequencing primers were the 17-mer universal M13 primer (Amersham) and 15- to 17-mer oligonucleotides synthesized to extend DNA sequence information beyond what was determined with M13 primers.

DNA sequence determination of the cDNA clone (Fig. 3A) showed that it was derived from a functionally rearranged β gene composed of the V β 5.1 leader, V β 8.2; a short portion of the D β 2 segment, J β 2.3; and the C β 2 gene segment. One of the two β alleles, isolated in clones λ 8 by restriction mapping and DNA sequencing showed that it contains the rearranged V β 8.2 gene about 100bp downstream of an apparently inactive V β 8.2 leader exon, while the V β 5.1 leader sequence is located about 2.5kb farther upstream (Fig. 3B). Splicing of the V β 5.1 leader exon to the V β 8.2 exon in T cells derived from C57BL/6 mice has been described by Chou, et al. (1987) Proc. Natl. Acad. Sci. USA 84, 1992-1996. The second β allele in B6.2.16 contains an incomplete D β 1-J β 2.5 rearrangement (not shown).

For the production of transgenic mice, a functionally rearranged β gene with long 5' and 3' flanking sequences was used. This was because initial experiments with clone λ 8 containing only 1.2kb of 5' and 3kb of 3' flanking sequence failed to give

expression in nine independently obtained transgenic mice.

Transgene Reconstitution

From the genomic clone λ 9, a 13 kb SaII-SacII
5 fragment, containing the functionally rearranged V β
gene with 9 kb of 5' flanking sequence, was fused by
way of a unique Sac II site to a 23 kb Sac II-Sal I
fragment containing J β 2 and C β 2 gene segments
together with 18kb of 3' flanking sequence. The
10 latter fragment was isolated from cosmid clone cos H-
2^dII-1.14T derived from a BALB/c liver library,
previously described (Steinmetz, et al. (1986) Cell
44, 895-904). See Fig. 3. Both fragments were
ligated in the presence of Sal I-digested pTCF cosmid
15 vector DNA arms. Ligation was checked by agarose gel
electrophoresis. In vitro packaging and
transformation of E. coli strain 490A was carried out
as previously described (Steinmetz, et al. (1985)
Methods in Molecular Immunology in Immunological
20 Methods, Vol. III, Letkovits and Pernis, eds.
(Orlando, Fla., Academic Press)). Several clones
were picked and clone cos HY β 9-1.14-5, containing the
reconstituted transgene as shown in Fig. 3B, was
identified by restriction endonuclease mapping.

25

Transgenic Mice

The 36 kb insert of cos HY β 9-1.14-5 was released by
SaII digestion and isolated by preparative agarose
gel electrophoresis and electroelution. The DNA was
extracted twice with phenol-chloroform and
precipitated with ethanol. The DNA pellet was
30 dissolved in ultrapure water and dialyzed against 10
mM Tris-HCl, 0.1 mM EDTA (pH 7.4). The DNA was
adjusted to a final concentration of 4 μ g/ml.
Fertilized mouse eggs were recovered in cumulus from

the oviducts of superovulated (CBA/BrA x C57BL/LiA)F₁ females that had mated with F₁ males several hours earlier. Strains CBA/BrA and C57BL/L.A were obtained from The Netherlands Cancer Institute breeding facility, Amsterdam. Approximately 100 copies of the β gene were injected in the most accessible pronucleus of each fertilized egg, as described by Hogan, et al. (1986) Manipulating the Mouse Embryo (Cold Spring Harbor, New York, Cold Spring Harbor Laboratory Press. Microinjected eggs were implanted into the oviducts of one-day pseudopregnant (CBA/BrA x C57BL/LiA)F₁ foster mothers and carried to term. Several weeks after birth, total genomic DNA was isolated from tail biopsies of the pups. Mice that had incorporated the injected DNA in their genome were used for further breeding.

Southern and Northern Blot Analyses

Tail DNA was purified from the terminal quarter of tails of 4-week-old mice. The skin was separated from the bone and homogenized in 1ml of 1% NaCl, 10mM EDTA (pH 8.0), on ice by using a Polytron with a PTA7 blade. DNA was isolated by phenol-chloroform extraction and dialyzed against 10mM Tris, 1mM EDTA (pH 8.0) (TE). One milliliter of 6% p-aminosalicylate was added to the first phenol extraction. Mouse organs were chopped in ice-cold PBS and homogenized in 4M guanidinium isothiocyanate. 0.5mM sodium citrate (pH 7.0), 0.1M β -mercaptoethanol, 0.5% Sarkosyl, using a Polytron as above. From the homogenate, RNA and DNA were isolated by differential centrifugation as described previously (Maniatis, et al. (1982) Molecular Cloning: A Laboratory Manual (Cold Spring Harbor, New York, Cold Spring Harbor Laboratory Press). DNA-containing fractions were dialyzed against TE buffer

-36-

for 6 hr. extracted with phenol-chloroform three times, and dialyzed against TE overnight.

Total cellular RNA (5 to 10 μ g) was separated on 1.5% formaldehyde-agarose gels according to Rave, et al. (1979) Bucl. Acad. Res. 6, 3559-3527) and transferred to Zeta probe membranes (BioRad) or BA85 nitrocellulose filters (Schleicher & Schull). Genomic DNAs (10 μ g) were digested to completion with restriction endonuclease, separated on 0.6% agarose gels, and transferred to Zeta probe membranes according to Reed and Mann (1985) Nucl. Acad. Res. 13, 7201-7221.

Northern blot hybridization with labeled restriction fragments was performed by using the same conditions as for Southern blot hybridization described by Steinmetz (1986), supra. Hybridization with the 30-mer oligonucleotide containing the VDJ joining region of the transgene was done in 900mM NaCl, 90mM Tris-HCl (pH 7.2), 6mM EDTA, 10 x Denhardt's solution, 1% SDS, and 500 μ g/ml E. coli DNA at 42°C overnight. After hybridization the filter was washed twice in 6x SSC, 0.1% SDS, at room temperature and twice in 3x SSC, 0.1% SDS, at 60°C for 10 min. each. Southern blot hybridization was carried out as described by Uematsu, et al. (1988) Immunogenetics 27, 96-101. Filters were exposed with Kodak X-Omat S or X-Omat AR films at -70°C for 2 hr. to 1 month using intensifying screens.

For hybridization we used previously described C α and C β probes were used (Dembić, et al. (1985) Nature 314, 271-273; Snodgrass, et al. (1985) Nature 313, 592-595. The J β 2 probe is a 1.2kb C1aI-EcoRI fragment located immediately downstream of the J β 2

-37-

gene segments (Chien, et al. (1984) Nature 309, 322-326). The D β 1 probe is a 1.4kb upstream of the D β 2 gene segment (Siu, et al. (1984) supra. The C β 1-specific probe was derived from the 3' untranslated region as described by Gascoigne, et al. (1984) Nature 310, 387-391. The transgene-specific VDJ probe is an oligonucleotide of 30 residues in length identical to the complementary strand of the underlined cDNA sequence shown in Fig. 3A from position 581 to 610. V β 7, V β 9, V β 11, V β 12, V β 14, V β 15 and V β 16 probes have been described by Lai, et al. (1987) Proc. Natl. Acad. Sci. USA 81, 3846-3850.

Surface Staining of Lymphocytes

Single-cell suspensions were prepared from lymph node or spleen. Cells were incubated with various monoclonal antibodies purified from supernatants of antibody-producing hybridoma cells. Cells (106) were incubated with a saturating dose of antibody at 0°C for 20 min, washed, and incubated again with fluorescein isothiocyanate (FITC) coupled goat anti-mouse immunoglobulin antibodies. In some experiments, antibodies coupled with biotin were used in the first step followed by incubation with FITC-avidine (Kisielow, et al. (1984) J. Immunol. 133, 1117-1123).

Modulation and Immunoprecipitation of Surface Antigens

For modulation, cells were incubated with an excess of F23.1 antibodies (50 μ g/ml) and rabbit anti-mouse immunoglobulin antibodies for 24 hr. at 37°C. Viable cells were obtained by spinning the suspension through Ficoll. For immunoprecipitation, 10⁷ B6.2.16 cloned T cells, C57BL/6 T cells, and 93.2 T cells were labeled with 0.5 mCi of NaI (Amersham) by the

-38-

lactoperoxidase/glucoseoxidase method (Goding (1980) J. Immunol. 124, 2082-2088) for 20 min. at room temperature in PBS. The cells were washed five times with ice-cold PBS containing 0.1% NaN₃ and 17mg/ml KCI and were then lysed for 30 min. on ice with a buffer containing 2% Triton X-100, 20mM Tris (pH 8), 150mM NaCl, 3 mM MgCl, 0.1mM PMSF, and 5mM iodoacetamide. The lysate was centrifuged for 5 min. in an Eppendorf centrifuge. Half of each sample was precleared four times with 100 μ l of F23.1 antibodies coupled to Sepharose beads. Precleared and untreated lysates were then incubated with 5 μ g of an anti- β panspecific antibody (Traunecker, et al. (1986) Eur. J. Immunol. 16, 851-854) for 2 hr. on ice, and subsequently incubated with 35 μ l of 50% protein A beads. All samples were washed with high salt (lysis buffer containing 1 mg/ml ovalbumin and 0.65 M NaCl) and low salt (lysis buffer containing 0.15M NaCl) buffers. Samples were reduced with 2-mercaptoethanol and analyzed on a 12.5% polyacrylamide gel. Autoradiography was accomplished with Kodak X-Omat AR film and intensifying screens.

Generation of Cytolytic T cells and Cytolytic Assay

Spleen (10^7) or lymph node (10^7) cells were cultured with 10^7 x-irradiated (2000 rads) allogeneic stimulator cells for 5 days in 4ml of culture medium. ^{51}Cr -labeled target cells were prepared by stimulating spleen cells from various mouse strains with concanavalin A (Con A) (2.5 μ g/ml) for 48 hr. at 10^6 cells per ml. Con A blasts were purified by centrifugation over Ficoll, and viable cells were ^{51}Cr -labeled by incubation in ^{51}Cr for 1 hr. at 37°C. In the cytolytic assay, 10^4 ^{51}Cr -labeled targets were incubated with various numbers of cytolytic T cells

for 3.5 hr. at 37°C in serial bottom wells in 200 μ l of medium. The plates were centrifuged and 100ml of supernatant removed to determine released ^{51}Cr (Pohlitz, et al. (1979) Eur. J. Immunol. 9, 681-690).

- 5 For blocking of activity, cytolytic T cells were incubated for 30 min. at 37°C with F23.1 antibodies (50 μ g/ml). Then ^{51}Cr -labeled targets were added and the cytolytic assay were performed as described above.

10 Establishment of Cell Lines and
 T Cell Clones from Transgenic Mice

- Cell lines were made by stimulation of spleen cells with either Con A or allogeneic x-irradiated stimulator cells in medium containing IL-2. After 10
15 to 14 days, the cells were washed and restimulated (10⁶ responder cells, 10⁷ x-irradiated stimulator cells) in IL-2-containing medium. Viable cells were obtained by centrifugation over Ficoll. Cloning was carried out by seeding 0.3 cells per well in 96-well
20 microtiter plates containing 10⁶ X-irradiated feeder spleen cells per well and 200 μ l of IL-2-containing medium. After 1 to 2 weeks, growing colonies were transferred together with 10⁷ x-irradiated feeder cells into 2ml of IL-2-containing medium in 24-well
25 Costar plates. From then on, restimulations were carried out at 7-day intervals with 10⁶ cloned cells and 10 x-irradiated stimulators per 1ml of IL-2-containing culture medium.

30 Transgenic Mice Express the
 Introduced T Cell Receptor β Gene

Thirteen pups were born that contained from 1 to about 50 copies of the transgene, according to Southern blot analysis of tail DNA. Of these mice, seven were analyzed after splenectomy for

transcription with the oligonucleotide probe that covered the VDJ joining region and that was therefore specific for the transgene. Six of the seven mice showed a 1.3kb full-length transcript. As observed before for other transgenes, no simple correlation between copy number and transcriptional activity was seen. T lymphocytes obtained from lymph nodes of four transgenic mice were subsequently analyzed for cell-surface expression of the transgenic β chain on a fluorescence-activated cellsorter (FACS) by using F23.1 antibodies (Table 1). While about 10%-20% of T cells were F23.1-positive in mice 90 and 91, practically all of the T cells from mice 93 and 95 were stained with the F23.1 antibody. This indicates that most, if not all, of the T cells in mice 93 and 95 express the transgenic β chain on the cell surface. Since a background of about 10%-20% F23.1-positive T lymphocytes is expected for these mouse strains (Staers, et al. (1985) J. Immunol. 134, 3994-4000), it is unlikely that the transgenic β chain is expressed in mice 90 and 91. In agreement with the FACS analysis, Northern blot analysis did not show any transgene transcription in T lymphocytes of mouse 90 (Table 1).

-41-

Table 1

Transgenic Mice Express the
Introduced T Cell Receptor β Gene

<u>Founder</u>	<u>Off- spring</u>	<u>Sex^a</u>	<u>Copy Number^b</u>	<u>Transgene Tran- scription</u>	<u>T Cell Surface Expression^c</u>
90		m	1	-	10%-20%
91		f	20	ND	10%-20%
92		f	10	ND	ND
93		f	5	+	>98%
	93.1	m	0	ND	ND
	93.2	m	20	+	>98%
	93.4	m	2	+	>98%
	93.9	f	20	+	>98%
94		m	3	+	ND
95		f	2	+	>98%
	95.39	m	2	+	ND
	95.40	m	2	ND	ND

(a) Sex is indicated by m (male) and f (female).

(b) Copy number of the transgene was estimated from signal intensities on Southern blots with PvuII-digested tail DNA hybridized with the J β 2 probe.

(c) Surface expression of F23.1-positive β chains on lymph node T cells was quantitated by FACS analysis.

ND - Not determined.

The male founder mouse 93 was subsequently crossed with female C57L mice (a F23.1-negative mouse strain; Behlke, et al. (1986) Proc. Natl. Acad. Sci. USA **83**, 767-771. Eighteen offspring mice from the first generation were analyzed for the inheritance of the transgene by Southern blot hybridization. Three (1 male, 2 females) contained about 20 copies of the

transgene, 5 (4 males, 1 female) contained about 2 copies, and 10 lacked the transgene (Fig. 4). This blot shows the analysis of 9 mice. It demonstrates the inheritance of the transgene (indicated by the 7.7kb band) in either 2 (mice 93.3, 93.4, 93.7) or 20 (93.2, 93.9) copies. The 5.5kb band represents the endogenous β locus. Southern blot analysis of PvuII-digested tail DNA was carried out as described with a J β 2 probe. The segregation of the transgene into 2 and 20 copies could be due to integration of injected DNA into two unlinked chromosomal loci, or could reflect a deletion or amplification event after integration at a single site. Analysis of second generation offspring mice shows that the two different forms of the transgene are stably inherited. The transgene is therefore present in germ-line DNA and transmitted in a Mendelian fashion to both males and females. Mice with 2 or 20 copies of the transgene were analyzed for cell-surface expression with F23.1 antibodies. Lymph node T cells stimulated in vitro by irradiated allogeneic DBA/2 spleen cells were incubated with V β 8-specific F23.1 antibodies followed by FITC-labeled sheep anti-mouse immunoglobulin antibodies, and were analyzed on FACS. In parallel, the expression of CD3 molecules was monitored by using the 145.2C11 monoclonal antibody (Leo, et al. (1987) Proc. Natl. Acad. Sci. USA 84, 1374-137??) followed by FITC-labeled goat anti-hamster immunoglobulin antibody. As shown in Fig. 5, most, if not all, of the peripheral T lymphocytes analyzed in mice 93.2 (20 copies) and 93.4 (2 copies) stain with the V β 8-specific monoclonal antibody. The staining intensity is similar in both T lymphocyte populations, indicating that despite the difference in copy number, these mice express the transgene to the same extent.

Lymph node cells from mice with 2 copies of the transgene were analyzed for the ratio of CD4- and CD8-positive T cells. Lymph node cells from a transgenic male mouse were 45% CD4⁻ and 18% CD8⁺, and from a transgenic female mouse, 35% CD4⁺ and 29% CD8⁺; control cells from a C57BL/6 mouse were 45% CD4⁺ and 32% CD8⁺. These numbers do not deviate significantly from those normally seen. Thus, although the transgenic β chain, derived from a class I-restricted cytotoxic T cell clone, is expressed in most, if not all, of the T lymphocytes, it does not affect the normal ratio of CD4- and CD8-positive T cells.

Transcription of the Transgene
is Largely T Cell-Specific

Mouse 93.2, containing 20 copies of transgene was analyzed for tissue-specific expression of the transgene. Northern blot analysis using the VDJ oligonucleotide as a probe reveals strong transcription of the thymus, while little or not transcription is seen in the other tissues analyzed (Fig. 6A). Similar results were obtained for transgenic mice 93.4 and 95.39, containing 2 copies of the transgene (not shown). The small amounts of transcripts we see in some nonlymphoid tissues could be due to infiltrating T lymphocytes or incomplete inactivation of the transgene. N lymphocytes also show a low level of transgene transcription. Both the VDJ oligonucleotide and the C β probe identify β chain transcripts in B lymphocytes, while α chain transcripts, even after a longer exposure, are not seen (Fig. 6B). The observed transcripts of the transgene are therefore not due to residual T cell contamination of the B-cell preparation, but reflect

an about 20-fold lower level of transgene transcription in B as compared to T lymphocytes.

Transgenic T Lymphocytes Express Only
F23.1-Positive β Chains, Which Participate
in the Formation of Functional Receptors

5 The F23.1 staining experiments indicate that the transgene is expressed in the vast majority of T lymphocytes. It is possible that additional β chains are encoded by endogenous genes. Several independent
10 approaches were used to check this possibility. In a first kind of experiment, membrane extracts from T cells of transgenic mice were "precleared" of V β 8 proteins by using F23.1 antibodies coupled to Sepharose beads. As shown in Fig. 7, after
15 preclearing, no further β chains were precipitated by a panspecific antiserum (Traunecker, et al. (1986) Eur. J. Immunol. 16, 851-854) for mouse β chains. When the original male-specific clone B6.2.16 and T cells from C57BL/6 mice were analyzed for controls,
20 the panspecific antiserum precipitated β chains from the alter but not from the former cells after preclearing with F23.1 antibodies. In a second kind of experiment, T cell receptors were modulated (by capping and endocytosis) by F23.1 antibodies and
25 rabbit anti-mouse immunoglobulin at 37°C. Modulated and nonmodulated cells were then stained with anti-CD3 antibodies (Leo, et al. (1987) supra.). As shown in Fig. 8, modulation with F23.1 antibodies removed all T cell receptor-associated CD3 molecules from
30 cell surfaces of the transgenic mice and from the B6.2.16 clone but not from other T cells. A third kind of experiment showed that the activity of cytolytic T cells from transgenic mice on allogeneic DBA/2 and C3H/HeJ target cells was completely blocked
35 by F23.1 antibodies, whereas those from normal C57BL/6 mice were not inhibited significantly when

used on the same targets (Fig. 9). Lysis of allogeneic target cells after in vitro stimulation was specific. Cytolytic T cells from transgenic mice stimulated with DBA/2 cells lysed DBA/2 but not C3H/HeJ target cells, and stimulation with C3H/HeJ cells gave rise to C3H/HeJ-directed, but not DBA/2-directed, killer cells (not shown). Taken together, these experiments indicate that only β chains encoded by the transgene are expressed o T lymphocytes of the transgenic mice and that they are used to form functional T cell receptors.

Endogenous β Genes are Incompletely Rearranged

Rearrangement of endogenous β genes can be distinguished from the rearranged transgene by using specific hybridization probes. With a $C\beta$ probe, rearrangements of the $C\beta 1$ locus can be separated from those at the $C\beta 2$ locus (which is present in the transgene) if HindIII-digested T cell DNA is analyzed. This analysis clearly shows that the 9.4kb germ-line HindIII fragment containing the $C\beta 1$ locus is rearranged in transgenic T lymphocytes from mouse 93.2 (Fig. 10A). The rearrangements, however, appear to be qualitatively different from those seen in normal C57L T lymphocytes used as a control. While no discrete rearranged fragments appear in C57L T cell DNA, a series of fragments (not seen in a B cell control) is evident in T cell and Thymus DNA from transgenic mouse 93.2. Also, mouse 93.1, which did not inherit the transgene, looks like the C57L control.

With a probe derived from the 5' flanking region of the $D\beta 1$ gene segment, a series of discrete rearranged fragments is again seen in transgenic but not in control T cell, nor in transgenic B cell, DNA (Fig.

-46-

10B). This result shows that partial D β 1 rearrangements, presumably to J β 1 and J β 2 gene segments, are present in unusually high frequencies in the transgenic mouse. Indeed, the sizes of the
5 observed rearranged fragments are in agreement with this assumption. Since these fragments are identified with a 5' flanking probe of D β 1, they represent partial rearrangements that do not involve V β gene segments. Inversion of the V β 14 gene
10 segment, located downstream of C β , would not delete the 5' flanking sequence of D β gene segments (Malissen, et al. (1986) Nature 319. 28-33). Analysis of T cell clones from transgenic mice (see below) with a V β 14 probe, however, shows that
15 endogenous V β 14 genes have not been rearranged to D β gene segments. In agreement with the Southern blot analysis, a Northern blot analysis of thymus and peripheral T cell RNA with a mixture of ³²P-labeled V β 7, V β 9, V β 11, V β 12, V β 14, V β 15 and V β 16 gene
20 segments did not reveal V β gene transcripts in transgenic mouse 93.2, while a clear signal was obtained with a C57BL/6 control (not shown).

In contrast to the D β 1 gene segment, the D β 2 gene segment does not undergo frequent rearrangements in
25 transgenic mouse 93.2. As shown in Fig. 10C, it rather seems that most of the D β 2 gene segments remain in germ-line configuration in 93.2 thymocytes and T cells. Mouse 93.4, containing only 2 rather than 20 copies of the transgene, shows a similar high
30 frequency of partial D β 1 rearrangements (not shown). Furthermore, mouse 95.39, independently generated, and a transgenic mouse obtained by coinjection of the same β gene together with a T cell receptor α gene show a predominance of partial D β 1 rearrangements
35 (not shown). Thus neither the high copy number of the

-47-

transgene in mouse 93.2 not its peculiar chromosomal location causes this unusual pattern of rearrangement.

5 The results presented so far do not exclude a low but
still significant number of complete endogenous VDJ
rearrangements. To obtain a more quantitative
estimate, we analyzed nine T cell clones obtained
from mice 93.2 and 93.4 with the same C β , D β 1, and
10 D β 2 probes. Of the 18 endogenous β loci screened,
13 rearranged D β 1 to J β 1, 2 fused D β 1 with J β 2, 3
joined D β 2 with J β 2, and 2 show unusual
rearrangements of both D β 1 and D β 2 gene segments that
were not further characterized (Table 2). No
complete VDJ rearrangements were found, indicating
15 that they occur very rarely if at all.

-48-

Table 2
T Cell Clones Analyzed

<u>Mouse</u>	<u>Clone^b</u>	<u>Rearrangements^a</u>			
		<u>Dβ1 to Jβ1</u>	<u>Dβ1 to Jβ2</u>	<u>Dβ2 to Jβ1</u>	<u>Dβ1 to Jβ1^c</u>
93.2	2	1		1	1
	8	1	1		
	14	1			1
	16	2 ^d			
	17	2			
	18	1	1		
	19	2			
	20 ^e	1		2	
	2	2			
93.4	2	2			

Notes:

- (a) Rearrangements of the two homologous loci in each T cell clone were analyzed with C β , D β 1, and D β 2 probes.
- (b) T cell clones were obtained as described in Experimental Procedures.
- (c) The sizes of these rearranged fragments do not correspond to those expected for DJ rearrangements.
- (d) Indicates that DJ rearrangements have occurred on both homologous chromosomes.
- (e) Clone 20 contains one β locus in germ-line configuration.

Expression of the Transgene Regulates
Rearrangement of Endogenous β Genes

The majority of endogenous β genes in the transgenic mice show rearrangements of D β 1 to J β 1 gene segments. No endogenous VDJ rearrangements are seen. The

preponderance of incomplete D β 1 to J β 1 rearrangements is highly unusual, and is independent of the copy number of the transgene and its chromosomal location.

5 It can be deduced from the analysis of T cells and T cell clones that approximately half of the nonproductive β rearrangements in normal T cells represent VDJ rearrangements. Of eight nonfunctional alleles that have been characterized in T cell clones, five show partial DJ rearrangements (at least
10 three of which are D β 1-J β 2) and three show aberrant VDJ rearrangements. Expression of the introduced β gene in transgenic mice therefore blocks endogenous β gene rearrangement between D \rightarrow J and V \rightarrow DJ joining.

Example 2

15 T CELL DEPLETED TRANSGENIC MICE

Transgenic mice depleted in T cell population were produced by introducing a transgene encoding the functionally rearranged T cell β receptor chain of Example 1 wherein approximately 90% of the variable
20 region of the β receptor gene was deleted.

Transgene Constructions

The 20kbp KpnI fragment derived from cosHY β 9-1.14-5 (Fig. 3B) was subcloned in PUC18. This plasmid was deposited with the Centraalbureau Voor Schimmelcultures (CBS) Baarn, The Netherlands on
25 December 5, 1988 and assigned deposit No. _____. The construction of the deletion transgenes from this plasmid is illustrated schematically in Figs. 11A, 11B and 12. The functional TCR β gene from the
30 B6.2.16 contains in addition to the fused V β 8.2, D β 2 and J β 2.3 segments the J segments J β 2.4, J β 2y and J β 2.6. A 4.0kb XbaI fragment contains the VDJ join and the remaining J segments. This fragment was

inserted into pSP64 (Promega, Madison, Wisconsin to form pSP4X). In this region of the KpnI fragment three PstI sites are present, the first at 17bp from the start of the V β 8.2 segment, the second between the J β 2.5 and the J β 2 ψ and the third at 22bp downstream from the J β 2 ψ segment. The 707bp located between the first and the last PstI sites were deleted from the KpnI fragment thereby removing all but the first six codons of the V β 8.2 segment, D β 2 and J β 2-J β 5. Two complementary oligonucleotide sequences were synthesized corresponding to the last six codons of the J β 2.3 segment and the 34bp region downstream from the J β 2.3 segment including the splice donor sequences. The oligomers were prepared in such a way that a PstI site is generated at the border in order to facilitate insertion into a PstI site. Digestion of pSP4X and religation resulted in the pSP2.7XP plasmid carrying a 2.7kbp XbaI/PstI fragment in which all of the V β 8.2 sequences, except for the first 17bp, and the 3' sequences up to the Xba site were deleted. pSP2.7XP was cleaved with PstI and religated in the presence of excess of reannealed oligomers (for sequence data see Fig. 11). The sequence chosen at the 5' and 3' of the oligomers was chosen such, that insertion of reannealed oligomer in the correct orientation restores only the 3' PstI restriction site. Plasmid pSP2.7XP contained one copy of the synthetic of 57mer in the correct orientation. In the PstI site of pSP2.7XP the 550bp PstI fragment containing the J β 2.6 segment derived from the pSP4X subclone was inserted to yield the pSP3.2X* clone. pSP3.2X* carries a 3.2kb XbaI fragment identical to 4.0 kbp XbaI insert of the pSP4X clone except for the deletion of the 707bp PstI fragment. Finally, the 4.0kb XbaI fragment in the 20kb KpnI fragment was replaced by the 3.2kb XbaI

insert of pSP3.2X* via multiple cloning steps involving the unique SacII site. Such multiple cloning steps are shown in Fig. 12. This resulted in the formation of the ΔV -TCR β λ ov ϵ .

5 A ΔV_{L} -TCR β clone containing one extra nucleotide was obtained in the same way, except that the pSP2.7XP subclone incorporated oligomers that contained the extra nucleotide in the coding region (Fig. 11). Because of the presence of the additional nucleotide
10 the mRNA transcribed from this construct can not be translated into a polypeptide carrying a C region. This construct served as a control after introduction into transgenic mice. All manipulations involving the deletion of the 707bp PstI fragment and the
15 insertion of the oligomers were checked by nucleotide sequence analysis.

Generation of Transgenic Mice

The DNA fragments (ΔV -TCR β or ΔV_{L} -TCR β) that were used for injection were released from the above
20 vectors using the appropriate restriction endonucleases and purified as described earlier (Krimpenfort, et al. (1988) Embo. J. 7, 745). The final DNA concentration was adjusted to 2 μ g/ml. Fertilized eggs were recovered in cumulus from the
25 oviducts of superovulated (CBA/BrA X C57Bl/LiA) F1 females that had mated with F1 males several hours earlier. The DNA fragments were injected into the most accessible pronucleus of each fertilized egg essentially as described (Hogan, B.J.M., et al.
30 (1985) Manipulation of the Mouse Embryo: A Laboratory Manual, Cold Springs Harbor Laboratory Press, Cold Spring Harbor, N.Y.). After overnight culturing two-cell-stage embryos were implanted into the oviducts of pseudopregnant fosters (F1 or BDF)

and carried to term. Several weeks after birth total genomic DNA was prepared from tail biopsies as described (Hogan, et al., supra)

DNA Analysis

5 For Southern blot analysis, 8 μ g total genomic DNA was digested with restriction endonucleases as recommended by the supplier, separated on agarose
10 gels and transferred to nitrocellulose. Filters were hybridized to 32 P-labeled probes as described (Cuypers, H.T., et al. (1984) Cell 37, 141). Final wash was at 0.1 X SSC, 42°C. The probe used for the transgene analysis was an EcoRI/HincII fragment located downstream from the J β 2 cluster. This probe recognizes a 6.0kbp PvuII fragment in germline DNA,
15 whereas a 7.9kbp PvuII fragment was diagnostic for the Δ V constructs.

RNA Analysis

5-15 μ g of total RNA, prepared by the LiCl-urea method (Auffray, C., et al. (1980) Eur. J. Biochem. 177,
20 303) was separated on 1% agarose formaldehyde gels (Maniatis, et al. (1982), supra.) and transferred to nitrocellulose. Probes used for RNA analysis were C β probe comprising a cDNA fragment representing the C β 2 region which also hybridizes with C β 1 transcripts
25 (Snodgrass, H., et al. (1985) Nature 313, 592) and an actin probe (Dodemont, H.J., et al. (1982) EMBO J. 1, 167). These probes were 32 P-labeled by nick-translation. Hybridization conditions were as described (Cuypers, et al. (1984) Cell 37, 141).
30 Final wash was at 0.1 X SSC and 60°C.

FACS Analysis

Analysis was performed on a FACS II analyzer (Becton and Dickinson) using established protocols (Scollay,

-53-

R., et al. (1983) Thymus 5, 245). Single cell suspensions were prepared from spleen in balanced salt solution containing 5% fetal calf serum and 0.5 mg/ml sodium azide. Cells were stained with saturating amounts of antibodies and counterstained with FITC-coupled rabbit anti-rat antibodies (Vesmel, W., et al. (1987) Leukemia 1 155). The origin and specificity of the antibodies were as follows: L3T4 (monoclonal 129-19 (Pierres, A., et al. (1984) J. Immunol. 132, 2775), Thy-1 (monoclonal SgA D2-2), Lyt-2 (monoclonal 53-6-7).

Transgenic Mice

Two transgenic founders were obtained carrying the ΔV_{α} -TCR β construct. In both transgenic mice multiple copies had integrated in head-to-tail configuration (data not shown). RNA was isolated from spleen of these mice to check transcription of the ΔV_{α} -TCR β transgene (see below). One ΔV_{α} -TCR β founder was sacrificed for obduction. No abnormalities were observed (normal thymus, spleen).

Using the ΔV -TCR β construct eight transgenic mice were generated. In most of these mice multiple copies had integrated. Mouse #1670 (male) did not produce offspring and was sacrificed for analysis. Strikingly, the thymus in this founder was strongly reduced in size. No abnormalities were recorded in other tissues. All transgenic offspring from another transgenic mouse (#1733) had an even more pronounced phenotype as mouse #1670: no thymus was present, whereas other tissues appeared normal. No abnormalities were observed in negative offspring from this transgenic line. Splenocytes from founder #1670 and from positive offspring from #1733 were analyzed by FACS.

-54-

Transgene Transcription

As discussed above, the ΔV_{α} -TCR β differs in only one nucleotide from the ΔV -TCR β transcripts and therefore does not encode a polypeptide chain containing a TCR constant region.

Two RNA species are transcribed from TCR β genes, 1.0kb and 1.3kb in size. Partially (DJ) rearranged β genes yield the 1.0kb transcripts, whereas the fully (VDJ) rearranged β genes give rise to the 1.3kb RNAs.

In thymus both TCR β transcripts are present abundantly (Snodgrass, et al. (1985) supra.). In spleen, however, the 1.0kb TCR β transcript is hardly detectable. Correct transcription and processing of the ΔV gene constructs also lead to a 1.0kb mRNA.

Northern blot analysis using probes specific for the C β region cannot discriminate between ΔV_{α} -TCR β transcripts and DJ transcripts. Therefore, RNA from spleen rather than thymus was chosen for the analysis of the ΔV_{α} -TCR β transgene expression. RNA was

isolated from spleen, from control mice, and from mice carrying the ΔV_{α} -TCR β construct. As can be concluded from Fig. 13 the endogenous TCR β gene expression is not affected in the ΔV_{α} -TCR β transgenic mice. In spleen from control and transgenic mice a

1.3kb band of approximately equal intensity can be detected with the C β probe. This indicates that complete VDJ rearrangements are not inhibited in the ΔV_{α} -TCR β transgenic mice. However, RNA from the transgenic spleens shows a high expresison of a 1.0kb

RNA species that is probably derived from the transgene ΔV_{α} -TCR β . The difference in expression level between the two transgenic strains supports this interpretation. We conclude from the RNA analysis, that the ΔV_{α} -TCR β and therefor probably

also the ΔV -TCR β transgenes give rise to stable transcripts.

Results of FACS Analysis

As it was impossible to isolate a sufficient number of
5 cells from the thymus of the transgenic mice carrying
the ΔV -TCR β construct, FACS analysis was only
performed on splenocytes. Splenocytes were isolated
from founder #1670 and from transgenic progeny of
lines #1673 and #1733. As a control cell
10 suspensions were prepared from the spleen of negative
littermates. Splenocytes were stained with anti-sera
specific for the B cell marker surface immunoglobulin
and the T cell markers Thy-1, L3T4 (T4) and Lyt-2
(T8). The FACS data on cell suspensions from
15 different transgenic mouse lines were essentially the
same. In the control cell suspension the B cell
specific anti-sera show a reactive and an unreactive
cell population, representing B and T cells,
respectively (Fig. 14). In contrast, in the
20 transgenic cell preparation (1670) all cells stain
with the B cell specific anti-sera. This indicates,
that all cells in the transgenic spleens are of B
cell origin. This observation is supported by the
staining patterns obtained with the T cell specific
25 anti-sera. Whereas in the control spleen
approximately 30% of the cells can be stained with
the Thy-1 anti-serum, only a minor fraction if any of
the transgenic splenocytes are Thy-1 positive. The
same observations were made by FACS using the L3T4
30 and Lyt-2 anti-sera. Apparently, T cells are absent
from the spleen of the ΔV -TCR β transgenic mice.

Conclusion

The RNA analysis of spleen of transgenic mice
carrying the ΔV -TCR β gene construct shows that the

introduced ΔV constructs are highly expressed. Moreover, the ΔV_{α} -TCR β transgenic mice show, that the transcriptional activity of the ΔV constructs per se does not inhibit the complete rearrangements of the endogenous β chain genes, since in the ΔV_{α} -TCR β transgenic mice the 1.3kb VDJ is expressed at similar levels as in control mice. The analysis of the ΔV -TCR β transgenic mice is still preliminary, e.g. at the moment no data on $\gamma\delta$ bearing T cells are available. A few consistent observations were made: no or only a rudimentary thymic structure was left in several of independent ΔV -TCR β transgenic mice lines; FACS analysis on splenocytes from these mice demonstrated that T cells were virtually absent. The absence of T cells in the ΔV -TCR β transgenic mice can be explained in several ways. Firstly, the ΔV transgenic protein itself might be lethal for T cells. This is unlikely, since a high expression of the structurally similar D_{μ} -protein is not harmful for A-MuLV transformed pre-B cells. Moreover, it was shown that pre-B cells with a high D_{μ} -protein expression behave just like B cells that express a complete Ig heavy chain: they proceed in the B cell differentiation pathway with rearrangements of the light chain genes (Reth, et al. (1985) supra.). Secondly, the mutant chain might interfere with T cell maturation. It has been suggested that during thymic maturation T cells are subject to both positive and negative selection processes. These include the stimulation of thymocytes that are able to recognize foreign antigen in association with self MHC as well as the elimination of self-reactive cells. This selection requires the interaction with MHC determinants. TCRs play an essential role in these interactions. In ΔV -TCR β transgenic mice the mutant TCR β probably prevents expression of the endogenous β

chain genes by inhibiting the complete VDJ rearrangements in the same way as the D μ protein blocks further rearrangements of the Ig heavy chain genes in A-MuLV transformed pre-B cells. Consequently, T cells in these mice do not express a complete β chain gene and therefore do not carry a TCR, that can functionally interact with MHC determinants. As a result all T cells die. Three mechanisms could account for the actual loss of T cells: (i) active deletion of T cells that bear a non-functional TCR; (ii) the deleted TCR transgene forms a receptor complex that is selfreactive and therefore is clonally deleted; (iii) T cells of the ΔV -TCR β transgenic mice are unable to respond to proliferative stimuli. The available data do not discriminate between these alternatives.

Having described the preferred embodiments of the present invention, it will appear to those ordinarily skilled in the art that various modifications may be made to the disclosed embodiments, and that such modifications are intended to be within the scope of the present invention.

-58-

WHAT IS CLAIMED IS:

1. A transgenic non-human animal having a phenotype characterized by the substantial depletion of a mature lymphocytic cell type otherwise naturally occurring in the species from which said transgenic animal is derived, said phenotype being conferred by a transgene contained in at least the precursor stem cell of said lymphocytic cell type, said transgene comprising a DNA sequence encoding a lymphatic polypeptide variant which inhibits formation of the mature form of said lymphocytic cell type.
2. The transgenic non-human animal of Claim 1 wherein said lymphatic polypeptide variant encoded by said transgene is expressed in a precursor to said mature lymphatic cell type.
3. The transgenic non-human animal of Claim 1 wherein said mature lymphocytic cell type is selected from the group consisting of mature T-cells and plasma cells.
4. The transgenic non-human animal of Claim 3 wherein said lymphocytic cell type is mature T-cell.
5. The transgenic non-human animal of Claim 4 wherein said transgene encodes a carboxy-terminal portion of a T-cell receptor β -chain.
6. The transgenic non-human animal of Claim 5 wherein said carboxy-terminal portion of said T-cell receptor β -chain encoded by said transgene comprises the constant region of said T-cell receptor β -chain.

7. The transgenic non-human animal of Claim 5 wherein said carboxy-terminal portion of said T-cell receptor β -chain encoded by said transgene comprises the constant region of said T-cell receptor β -chain, a portion of the variable region of said T-cell receptor β -chain and the signal sequence of said T-cell receptor β -chain.

8. The transgenic non-human animal of Claim 7 wherein said portion of said variable region comprises a C-terminal portion of the J-segment of said variable region.

9. The transgenic non-human animal of Claim 8 wherein said portion of said variable region further comprises an N-terminal portion of the V-segment of said variable region.

10. The transgenic non-human animal of Claim 4 wherein said transgene is derived from a naturally-occurring T-cell receptor β -chain by the deletion of DNA sequences encoding all or part of the variable region of said T-cell receptor β -chain.

11. The transgenic non-human animal of Claim 4 wherein said transgene is derived from a DNA sequence encoding a lymphatic polypeptide having a first functional domain capable of suppressing expression of cognate endogenous alleles and a second functional domain capable of inducing formation of said mature lymphocytic cell type, said transgene being derived by deleting from said DNA sequence at least part of the sequence encoding said second functional domain.

12. The transgenic non-human animal of Claim 4 wherein said non-human animal is mouse.

-60-

13. The transgenic mouse of Claim 12 wherein said transgene comprises ΔV -TCR β .

14. A transgene comprising a DNA sequence expressible in a transgenic non-human animal, said transgene comprising a transgenic DNA sequence derived from a DNA sequence encoding a lymphatic polypeptide having a first functional domain capable of suppressing the expression of cognate endogenous alleles in said transgenic animal and a second functional domain capable of inducing the maturation of the lymphatic cell type normally expressing said endogenous alleles, said transgenic DNA sequence being derived by deleting from said DNA sequence at least part of the sequence encoding said second functional domain.

15. The transgene of Claim 14 wherein said transgene is capable of suppressing the expressing cognate endogenous alleles and incapable of inducing maturation of a lymphatic cell type is a transgenic non-human animal when the transgenic polypeptide encoded by said transgene is expressed in said transgenic animal.

16. A transgene comprising an expressible DNA sequence encoding a lymphatic polypeptide variant comprising a carboxy-terminal portion of a T-cell receptor β -chain.

17. The transgene of Claim 15 wherein said carboxy-terminal portion of said T-cell receptor β -chain comprises the constant region of said T-cell receptor β -chain..

18. The transgene of Claim 16 wherein said carboxy-terminal portion of said T-cell receptor β -chain comprises the constant region of said T-cell receptor β -chain, a portion of the variable region of said T-cell receptor β -chain and the signal sequence of said T-cell receptor β -chain.
19. The transgene of Claim 18 wherein said portion of said variable region comprises a C-terminal portion of the J-segment of said variable region.
20. The transgene of Claim 19 wherein said portion of said variable region further comprises an N-terminal portion of the V-segment of said variable region.
21. A transgene comprising an expressible DNA sequence, said DNA sequence being derived from a naturally occurring T-cell receptor β -chain by the deletion of DNA sequences encoding at least part of the variable region of said T-cell receptor β -chain.
22. A transgene comprising ΔV -TCR β .
23. A transgene comprising a DNA sequence encoding a lymphatic polypeptide variant.
24. The transgene of claim 23 wherein said lymphatic polypeptide variant encoded by said DNA is capable of being expressed in a precursor of a mature lymphocytic cell type.
25. A method for producing a transgenic non-human animal having a phenotype characterized by the substantial depletion of a mature lymphocytic cell type otherwise naturally occurring in the species

from which said transgenic animal is derived, said method comprising:

5 introducing a transgene into an embryonal target cell of a non-human animal, said transgene comprising a DNA sequence encoding a lymphatic polypeptide variant which is capable of inhibiting formation of said mature lymphocytic cell type when contained in a precursor stem cell of said mature lymphocytic cell type;

10 transplanting the transgenic embryonal target cell formed thereby into a recipient female parent; and

 identifying at least one offspring having said phenotype.

15 26. The method of Claim 25 wherein the lymphatic polypeptide variant encoded by said DNA sequence is capable of being expressed in a precursor of said mature lymphocytic cell type.

20 27. The method of Claim 26 wherein said transgene comprises a transgenic DNA sequence derived from a DNA sequence encoding a lymphatic polypeptide having a first functional domain capable of suppressing the expression of cognate endogenous alleles in said transgenic animal and a second functional domain
25 capable of inducing the maturation of the lymphatic cell type normally expressing said endogenous alleles, said transgenic DNA sequence being derived by deleting from said DNA sequence at least part of the sequence encoding said second functional domain.

30 28. The method of Claim 25 wherein said mature lymphocytic cell type is selected from the group consisting of mature T-cells and plasma cells.

29. The method of Claim 28 wherein said lymphocytic cell type is mature T-cell.

5 30. The method of Claim 29 wherein said transgene encodes a carboxy-terminal portion of a T-cell receptor β -chain.

31. The method of Claim 30 wherein said carboxy-terminal portion of said T-cell receptor β -chain encoded by said transgene comprises the constant region of said T-cell receptor β -chain.

10 32. The method of Claim 30 wherein said carboxy-terminal portion of said T-cell receptor β -chain encoded by said transgene comprises the constant region of said T-cell receptor β -chain, a portion of the variable region of said T-cell receptor β -chain and the signal sequence of said T-cell receptor β -chain.

15 33. The method of Claim 32 wherein said portion of said variable region further comprises a C-terminal portion of the J-segment of said variable region.

20 34. The method of Claim 33 wherein said portion of said variable region further comprises an N-terminal portion of the V-segment of said variable region.

25 35. The method of Claim 29 wherein said transgene comprises a DNA sequence derived from a naturally occurring T-cell receptor β -chain by the deletion of DNA sequences encoding at least part of the variable region of said T-cell receptor β -chain.

36. The method of Claim 29 wherein said non-human animal is mouse.

37. The method of Claim 36 wherein said transgene comprises ΔV -TCR β .

38. The method of Claim 37 wherein said embryonal target cell is a mouse zygote and said introduction of said transgene into said zygote is by microinjection.

39. A transgenic non-human animal having a phenotype characterized by the substantial depletion of a mature lymphocytic cell type otherwise naturally occurring in the species from which said transgenic animal is derived.




Fig-1A





Fig-1B




Figure 2: Genomic organization of the T cell receptor loci in the mouse (not to scale): V, D, J and C region segments are indicated and transcriptional orientations (where known) are marked with arrows (from Davis and Bjorkman, 1988).



Southern Blot Analysis of Oligopop Mice Derived from a Cross of the Female Transgenic Mouse 93 with CS7L Males

Fig-4



Surface Expression of F23.1 Positive β Chains on T Cells from Transgenic and Normal Mice

Fig-5


100
 110
 120
 130
 140
 150
 160
 170
 180
 190
 200
 210
 220
 230
 240
 250
 260
 270
 280
 290
 300
 310
 320
 330
 340
 350
 360
 370
 380
 390
 400
 410
 420
 430
 440
 450
 460
 470
 480
 490
 500
 510
 520
 530
 540
 550
 560
 570
 580
 590
 600
 610
 620
 630
 640
 650
 660
 670
 680
 690
 700
 710
 720
 730
 740
 750
 760
 770
 780
 790
 800
 810
 820
 830
 840
 850
 860
 870
 880
 890
 900
 910
 920
 930
 940
 950
 960
 970
 980
 990
 1000
 1010
 1020
 1030
 1040
 1050
 1060
 1070
 1080
 1090
 1100
 1110
 1120
 1130
 1140
 1150
 1160
 1170
 1180
 1190
 1200
 1210
 1220
 1230
 1240
 1250
 1260
 1270
 1280
 1290
 1300
 1310
 1320
 1330
 1340
 1350
 1360
 1370
 1380
 1390
 1400
 1410
 1420
 1430
 1440
 1450
 1460
 1470
 1480
 1490
 1500
 1510
 1520
 1530
 1540
 1550
 1560
 1570
 1580
 1590
 1600
 1610
 1620
 1630
 1640
 1650
 1660
 1670
 1680
 1690
 1700
 1710
 1720
 1730
 1740
 1750
 1760
 1770
 1780
 1790
 1800
 1810
 1820
 1830
 1840
 1850
 1860
 1870
 1880
 1890
 1900
 1910
 1920
 1930
 1940
 1950
 1960
 1970
 1980
 1990
 2000
 2010
 2020
 2030
 2040
 2050
 2060
 2070
 2080
 2090
 2100
 2110
 2120
 2130
 2140
 2150
 2160
 2170
 2180
 2190
 2200
 2210
 2220
 2230
 2240
 2250
 2260
 2270
 2280
 2290
 2300
 2310
 2320
 2330
 2340
 2350
 2360
 2370
 2380
 2390
 2400
 2410
 2420
 2430
 2440
 2450
 2460
 2470
 2480
 2490
 2500
 2510
 2520
 2530
 2540
 2550
 2560
 2570
 2580
 2590
 2600
 2610
 2620
 2630
 2640
 2650
 2660
 2670
 2680
 2690
 2700
 2710
 2720
 2730
 2740
 2750
 2760
 2770
 2780
 2790
 2800
 2810
 2820
 2830
 2840
 2850
 2860
 2870
 2880
 2890
 2900
 2910
 2920
 2930
 2940
 2950
 2960
 2970
 2980
 2990
 3000
 3010
 3020
 3030
 3040
 3050
 3060
 3070
 3080
 3090
 3100
 3110
 3120
 3130
 3140
 3150
 3160
 3170
 3180
 3190
 3200
 3210
 3220
 3230
 3240
 3250
 3260
 3270
 3280
 3290
 3300
 3310
 3320
 3330
 3340
 3350
 3360
 3370
 3380
 3390
 3400
 3410
 3420
 3430
 3440
 3450
 3460
 3470
 3480
 3490
 3500
 3510
 3520
 3530
 3540
 3550
 3560
 3570
 3580
 3590
 3600
 3610
 3620
 3630
 3640
 3650
 3660
 3670
 3680
 3690
 3700
 3710
 3720
 3730
 3740
 3750
 3760
 3770
 3780
 3790
 3800
 3810
 3820
 3830
 3840
 3850
 3860
 3870
 3880
 3890
 3900
 3910
 3920
 3930
 3940
 3950
 3960
 3970
 3980
 3990
 4000
 4010
 4020
 4030
 4040
 4050
 4060
 4070
 4080
 4090
 4100
 4110
 4120
 4130
 4140
 4150
 4160
 4170
 4180
 4190
 4200
 4210
 4220
 4230
 4240
 4250
 4260
 4270
 4280
 4290
 4300
 4310
 4320
 4330
 4340
 4350
 4360
 4370
 4380
 4390
 4400
 4410
 4420
 4430
 4440
 4450
 4460
 4470
 4480
 4490
 4500
 4510
 4520
 4530
 4540
 4550
 4560
 4570
 4580
 4590
 4600
 4610
 4620
 4630
 4640
 4650
 4660
 4670
 4680
 4690
 4700
 4710
 4720
 4730
 4740
 4750
 4760
 4770
 4780
 4790
 4800
 4810
 4820
 4830
 4840
 4850
 4860
 4870
 4880
 4890
 4900
 4910
 4920
 4930
 4940
 4950
 4960
 4970
 4980
 4990
 5000
 5010
 5020
 5030
 5040
 5050
 5060
 5070
 5080
 5090
 5100
 5110
 5120
 5130
 5140
 5150
 5160
 5170
 5180
 5190
 5200
 5210
 5220
 5230
 5240
 5250
 5260
 5270
 5280
 5290
 5300
 5310
 5320
 5330
 5340
 5350
 5360
 5370
 5380
 5390
 5400
 5410
 5420
 5430
 5440
 5450
 5460
 5470
 5480
 5490
 5500
 5510
 5520
 5530
 5540
 5550
 5560
 5570
 5580
 5590
 5600
 5610
 5620
 5630
 5640
 5650
 5660
 5670
 5680
 5690
 5700
 5710
 5720
 5730
 5740
 5750
 5760
 5770
 5780
 5790
 5800
 5810
 5820
 5830
 5840
 5850
 5860
 5870
 5880
 5890
 5900
 5910
 5920
 5930
 5940
 5950
 5960
 5970
 5980
 5990
 6000
 6010
 6020
 6030
 6040
 6050
 6060
 6070
 6080
 6090
 6100
 6110
 6120
 6130
 6140
 6150
 6160
 6170
 6180
 6190
 6200
 6210
 6220
 6230
 6240
 6250
 6260
 6270
 6280
 6290
 6300
 6310
 6320
 6330
 6340
 6350
 6360
 6370
 6380
 6390
 6400
 6410
 6420
 6430
 6440
 6450
 6460
 6470
 6480
 6490
 6500
 6510
 6520
 6530
 6540
 6550
 6560
 6570
 6580
 6590
 6600
 6610
 6620
 6630
 6640
 6650
 6660
 6670
 6680
 6690
 6700
 6710
 6720
 6730
 6740
 6750
 6760
 6770
 6780
 6790
 6800
 6810
 6820
 6830
 6840
 6850
 6860
 6870
 6880
 6890
 6900
 6910
 6920
 6930
 6940
 6950
 6960
 6970
 6980
 6990
 7000
 7010
 7020
 7030
 7040
 7050
 7060
 7070
 7080
 7090
 7100
 7110
 7120
 7130
 7140
 7150
 7160
 7170
 7180
 7190
 7200
 7210
 7220
 7230
 7240
 7250
 7260
 7270
 7280
 7290
 7300
 7310
 7320
 7330
 7340
 7350
 7360
 7370
 7380
 7390
 7400
 7410
 7420
 7430
 7440
 7450
 7460
 7470
 7480
 7490
 7500
 7510
 7520
 7530
 7540
 7550
 7560
 7570
 7580
 7590
 7600
 7610
 7620
 7630
 7640
 7650
 7660
 7670
 7680
 7690
 7700
 7710
 7720
 7730
 7740
 7750
 7760
 7770
 7780
 7790
 7800
 7810
 7820
 7830
 7840
 7850
 7860
 7870
 7880
 7890
 7900
 7910
 7920
 7930
 7940
 7950
 7960
 7970
 7980
 7990
 8000
 8010
 8020
 8030
 8040
 8050
 8060
 8070
 8080
 8090
 8100
 8110
 8120
 8130
 8140
 8150
 8160
 8170
 8180
 8190
 8200
 8210
 8220
 8230
 8240
 8250
 8260
 8270
 8280
 8290
 8300
 8310
 8320
 8330
 8340
 8350
 8360
 8370
 8380
 8390
 8400
 8410
 8420
 8430
 8440
 8450
 8460
 8470
 8480
 8490
 8500
 8510
 8520
 8530
 8540
 8550
 8560
 8570
 8580
 8590
 8600
 8610
 8620
 8630
 8640
 8650
 8660
 8670
 8680
 8690
 8700
 8710
 8720
 8730
 8740
 8750
 8760
 8770
 8780
 8790
 8800
 8810
 8820
 8830
 8840
 8850
 8860
 8870
 8880
 8890
 8900
 8910
 8920
 8930
 8940
 8950
 8960
 8970
 8980
 8990
 9000
 9010
 9020
 9030
 9040
 9050
 9060
 9070
 9080
 9090
 9100
 9110
 9120
 9130
 9140
 9150
 9160
 9170
 9180
 9190
 9200
 9210
 9220
 9230
 9240
 9250
 9260
 9270
 9280
 9290
 9300
 9310
 9320
 9330
 9340
 9350
 9360
 9370
 9380
 9390
 9400
 9410
 9420
 9430
 9440
 9450
 9460
 9470
 9480
 9490
 9500
 9510
 9520
 9530
 9540
 9550
 9560
 9570
 9580
 9590
 9600
 9610
 9620
 9630
 9640
 9650
 9660
 9670
 9680
 9690
 9700
 9710
 9720
 9730
 9740
 9750
 9760
 9770
 9780
 9790
 9800
 9810
 9820
 9830
 9840
 9850
 9860
 9870
 9880
 9890
 9900
 9910
 9920
 9930
 9940
 9950
 9960
 9970
 9980
 9990
 10000
 10010
 10020
 10030
 10040
 10050
 10060
 10070
 10080
 10090
 10100
 10110
 10120
 10130
 10140
 10150
 10160
 10170
 10180
 10190
 10200
 10210
 10220
 10230
 10240
 10250
 10260
 10270
 10280
 10290
 10300
 10310
 10320
 10330
 10340
 10350
 10360
 10370
 10380
 10390
 10400
 10410
 10420
 10430
 10440
 10450
 10460
 10470
 10480
 10490
 10500
 10510
 10520
 10530
 10540
 10550
 10560
 10570
 10580
 10590
 10600
 10610
 10620
 10630
 10640
 10650
 10660
 10670
 10680
 10690
 10700
 10710
 10720
 10730
 10740
 10750
 10760
 10770
 10780
 10790
 10800
 10810
 10820
 10830
 10840
 10850
 10860
 10870
 10880
 10890
 10900
 10910
 10920
 10930
 10940
 10950
 10960
 10970
 10980
 10990
 11000
 11010
 11020
 11030
 11040
 11050
 11060
 11070
 11080
 11090
 11100
 11110
 11120
 11130
 11140
 11150
 11160
 11170
 11180
 11190
 11200
 11210
 11220
 11230
 11240
 11250
 11260
 11270
 11280
 11290
 11300
 11310
 11320
 11330
 11340
 11350
 11360
 11370
 11380
 11390
 11400
 11410
 11420
 11430
 11440
 11450
 11460
 11470
 11480
 11490
 11500
 11510
 11520
 11530
 11540
 11550
 11560
 11570
 11580
 11590
 11600
 11610
 11620
 11630
 11640
 11650
 11660
 11670
 11680
 11690
 11700
 11710
 11720
 11730
 11740
 11750
 11760
 11770
 11780
 11790
 11800
 11810
 11820
 11830
 11840
 11850
 11860
 11870
 11880
 11890
 11900
 11910
 11920
 11930
 11940
 11950
 11960
 11970
 11980
 11990
 12000
 12010
 12020
 12030
 12040
 12050
 12060
 12070
 12080
 12090
 12100
 12110
 12120
 12130
 12140
 12150
 12160
 12170
 12180
 12190
 12200
 12210
 12220
 12230
 12240
 12250
 12260
 12270
 12280
 12290
 12300
 12310
 12320
 12330
 12340
 12350
 12360
 12370
 12380
 12390
 12400
 12410
 12420
 12430
 12440
 12450
 12460
 12470
 12480
 12490
 12500
 12510
 12520
 12530
 12540
 12550
 12560
 12570
 12580
 12590
 12600
 12610
 12620
 12630
 12640
 12650
 12660
 12670
 12680
 12690
 12700
 12710
 12720
 12730
 12740
 12750
 12760
 12770
 12780
 12790
 12800
 12810
 12820
 12830
 12840
 12850
 12860
 12870
 12880
 12890
 12900
 12910
 12920
 12930
 1294

Cells	B6.216	C57BL/6	93.2
pre-culture	-	-	-
anti-IL-1	-	-	-
F231	-	-	-




Gel Electrophoretic Analysis of Immunoprecipitated B Chains from Transgenic T Lymphocytes before and after Pre-culture with F231 Monoclonal Antibodies

Fig-7




Modulation of Surface Expression of CD3 Molecules on Vβ8+ T Cells by Vβ8 Specific, F231 Monoclonal Antibodies

Fig-8



Inhibition of Cytolytic Activity of Transgenic T Cells by Vβ8 Specific F231 Monoclonal Antibodies

Fig-9




Endogenous B Genes in Transgenic Mice: 93.2 Shows Predominantly Partial Dβ1, Jβ1 Rearrangements


Fig-10

a

4/7



b



Structure of the mutant TCRβ transgenes. Panel a: flowsheet of the construction of the transgenes; the KpnI fragment derived from cosHYβ9-1.14-5 served as the starting material for the construction; for details see sections results and materials and methods; relevant restriction enzymes: P: PstI, X: XbaI, K: KpnI. Panel b: representation in detail of the region mutated in the TCRβ chain encoding the variable region; the sequences inserted by the oligomers in the ΔV constructs are boxed; the additional nucleotide in the ΔV-TCRβ construct is indicated by an arrow.

F, 8 11

5/7






Fig-12



Surface marker staining profile of splenocytes from transgenic mouse #1670 and a control mouse (both 8 weeks of age). Single cell suspensions from spleen were stained with anti sera as indicated and evaluated by flow cytometry

Fig-14



Expression of the $\Delta V\alpha$ -TCR β transgene. Total RNA was isolated from the spleen of $\Delta V\alpha$ -TCR β transgenic mice (#1641 and #1642) and from the thymus and the spleen from a control mouse (C), electrophoresed on formaldehyde gels and transferred to nitrocellulose filters. Filters were hybridized to a probe specific for the C β constant region (C β) and a probe specific for actin (Act) to as a measure for the quantity and integrity of the RNA

Fig-13

INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 89/05210-1

I. CLASSIFICATION OF SUBJECT MATTER (if several classification symbols apply, indicate all) ⁶		
According to International Patent Classification (IPC) or to both National Classification and IPC IPC (5): C12N 5/00, C12N 15/00, C07H 21/04		
U.S. Cl. : 800/2 536/27		
II. FIELDS SEARCHED		
Minimum Documentation Searched ⁷		
Classification System	Classification Symbols	
U.S.	536/27 435/320, 317.1, 172.3 800/2	
Documentation Searched other than Minimum Documentation to the Extent that such Documents are Included in the Fields Searched ⁸		
Databases: DIALOG (Files 55(1981-1989), 311(1982-1986), 312(1987-1989), 154(1983-1989)), USPTO Automated Patent System (File USPAT, 1975-1989). See attachment for search terms.		
III. DOCUMENTS CONSIDERED TO BE RELEVANT ⁹		
Category [*]	Citation of Document, ¹¹ with indication, where appropriate, of the relevant passages ¹²	Relevant to Claim No. ¹³
X,P	Journal of Cellular Biochemistry (New York, USA), Volume: supplement 0(13 Part B), Issued January 1989, Krimpenfort et al., "Dominant negative mutation in TCR β chain gene causes T cell depletion in transgenic mice", page 181, abstract number F214, see the entire abstract.	1-39
Y,P	Proceedings of the National Academy of Sciences (Washington, USA); Volume 85, Issued December 1988, Boehmer et al., "Early expression of a T-cell receptor B-chain transgene suppresses rearrangement of the Vr4 gene segment", pages 9729-9732, see the entire document.	1-39
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>[*] Special categories of cited documents: ¹⁰</p> <p>"A" document defining the general state of the art which is not considered to be of particular relevance</p> <p>"E" earlier document but published on or after the international filing date</p> <p>"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>"O" document referring to an oral disclosure, use, exhibition or other means</p> <p>"P" document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step</p> <p>"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>"&" document member of the same patent family</p> </div> </div>		
IV. CERTIFICATION		
Date of the Actual Completion of the International Search		Date of Mailing of this International Search Report
14 February 1990		14 MAR 1990
International Searching Authority		Signature of Authorized Officer
ISA/US		<i>Jasmin C. Chambers</i> Jasmin C. Chambers

III. DOCUMENTS CONSIDERED TO BE RELEVANT (CONTINUED FROM THE SECOND SHEET)		
Category *	Citation of Document, with indication, where appropriate, of the relevant passages	Relevant to Claim No
Y	Nature (London, UK), Volume 333, Issued June 1988, Kisielow et al., "Tolerance in T-cell receptor transgenic mice involves deletion of nonmature CD4 ⁺ 8 ⁺ thymocytes", pages 742-746, see the entire document.	1-39
Y	Nature (London, UK), Volume 334, Issued July 1988, Bluthmann et al., "T-cell-specific deletion of T-cell receptor transgenes allows functional rearrangement of endogenous α - and B- genes", pages 156-159, see the entire document.	1-39
Y	Science (Washington, USA), Volume 241, Issued August 1988, Fenton et al., "Isotypic exclusion of $\gamma\delta$ T cell receptors in transgenic mice bearing a rearranged β -chain gene", pages 1089-1092, see the entire document.	1-39
Y	Science, (Washington, USA) Volume 236, Issued May 1987, Nussenzweig et al., "Allelic exclusion in transgenic mice that express the membrane form of immunoglobulin μ ", pages 816-819, see the entire document.	1-39
Y	Journal of Experimental Medicine (New York, USA), Volume 167, Issued June 1988, Nussenzweig et al., "Allelic exclusion in transgenic mice carrying mutant human IgM genes", pages 1969-1974, see the entire document.	1-39
Y	Proceedings of the National Academy of Science (Washington, USA), Volume 83, Issued February 1986, Behtke et al., "Murine T-cell receptor mutants with deletions of β -chain variable region genes", pages 767-771; see the entire document.	1-39

FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET

Y

Cell (Cambridge, USA), Volume 52,
 Issued March 1988, Uematsu et al.,
 "In transgenic mice the introduced
 functional T cell receptor B gene
 prevents expression of endogenous B
 genes", pages 831-841, see the
 entire document.

1-39

V. OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE¹

This international search report has not been established in respect of certain claims under Article 17(2) (a) for the following reasons:

1. ☐ Claim numbers _____, because they relate to subject matter¹² not required to be searched by this Authority, namely:

2. ☐ Claim numbers _____, because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out¹³, specifically:

3. ☐ Claim numbers _____, because they are dependent claims not drafted in accordance with the second and third sentences of PCT Rule 6.4(a).

VI. ☒ OBSERVATIONS WHERE UNITY OF INVENTION IS LACKING²

This International Searching Authority found multiple inventions in this international application as follows:

I. Claims 1-13 and 25-39, drawn to a transgenic non-human animal and a method of making the same, classified in Class 800, subclass 2.

II. Claims 14-24, drawn to a transgene, classified in Classes 536, subclass 27.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims of the international application.

2. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims of the international application for which fees were paid, specifically claims:

3. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claim numbers:

4. ☐ As all searchable claims could be searched without effort justifying an additional fee, the International Searching Authority did not invite payment of any additional fee.

Remark on Protest

☐ The additional search fees were accompanied by applicant's protest.

☐ No protest accompanied the payment of additional search fees.

Attachment to PCT/ISA/210, Part II.

II. FIELDS SEARCHED SEARCH TERMS:

transgenic, allelic exclusion, TCR, T-cell,
receptor, maturation, mutants, inventors' names.

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.